

**NOVEL COMPOSITIONS DERIVED FROM CRANBERRY AND
GRAPEFRUIT AND THERAPEUTIC USES THEREFOR**

Related Information

5 This application claims priority to U.S. Provisional Application No. 60/196,886,
entitled "NOVEL COMPOSITIONS DERIVED FROM CRANBERRY AND
GRAPEFRUIT AND THERAPEUTIC USES THEREFOR," filed on April 13, 2000,
incorporated herein in its entirety by this reference. The contents of all patents, patent
applications, and references cited throughout this specification are hereby incorporated
10 by reference in their entireties.

Background of the Invention

For millennia, humankind has relied on plant derivatives for the prevention and
treatment of a wide variety of ailments. For example, in China, various teas have been
15 used as a crude medicine for over 4,000 years. And more recently, there has been
considerable interest in taking advantage of various plant extracts as a source of health
promoting substances such as, natural oxidants, phenolic compounds, flavonoids,
tocochromanols, and beneficial fatty acids. In part, this trend is due to a growing body
of evidence demonstrating that some of these compounds have beneficial properties that
20 may be advantageous in preventing or delaying the onset of disease.

Indeed, several epidemiological studies considering the affect of diet on disease
such as, *e.g.*, cancer and hypercholesterolemia, have provided leads in the search for
naturally-occurring anti-cancer or anti-cholesterol agents. For example, some studies
suggest that plant-based diets, rich in whole grains, legumes, fruits and vegetables, may
25 reduce the risk of various types of cancer, including breast cancer (Steinmetz & Potter,
1991).

Similarly, other studies indicate that populations consuming large amounts of
cereal grains have lower cholesterol levels and a lower incidence of cardiovascular
disease. These studies have attributed these beneficial properties of cereal diets on the
30 presence of naturally occurring tocochromanols, and these compounds have been found
in a wide variety of plant sources (Quereshi *et al.*, *Am. J. Clin. Nutr.*, 53:1021S-6S
(1991)).

003344 0430
FOOTNOTES

Moreover, additional studies suggest that fruit products are a source of a number of health promoting phytochemicals (Johns *et al.*, *Recent Advances in Phytochemistry*, pp.31-52, Plenum Press (1997)).

Given that cancer and cholesterol-related diseases (*e.g.*, arteriosclerosis) are two
5 of the major causes of death in the United States, additional research on and
identification of fruit-derived therapeutic compounds which, for example, are useful in
treating or preventing such diseases, would be of great benefit.

Summary of the Invention

10 The present invention provides novel compounds and therapeutic compositions
(*i.e.*, formulations) derived from fruits, particularly cranberry and grapefruit, as well as
novel uses for the compounds and compositions. In particular embodiments, the
compounds are formulated as a pharmaceutical, a foodstuff (*e.g.*, added to a foodstuff to
enhance its nutritional and/or medical value), or a dietary supplement. In all cases, the
15 compounds and compositions contain, or are enriched for, health promoting components
(*e.g.*, phenolics, flavonoids, tocochromanols) that are useful in treating or preventing a
variety of health-related disorders and diseases.

Accordingly, in another embodiment, the present invention provides a method
for treating or preventing a disease in a subject, particularly a malignancy (*e.g.*, a
20 cancer), by administering to the subject (*e.g.*, orally or by injection) a therapeutically-
effective amount of a compound or composition of the invention. The malignancy can
be, for example, metastatic, breast cancer, or metastatic breast cancer. In another
embodiment, the present invention provides a method for treating or preventing
hypercholesterolemia in a subject, by administering to the subject a therapeutically-
25 effective amount of a compound or composition of the invention. The
hypercholesterolemia can be measured or detected by, for example, altered apoB levels.

Novel compositions of the invention are derived (*e.g.*, isolated) from, or contain
components of citrus or cranberry fruits. A preferred citrus fruit for use in the invention
is grapefruit, including pink grapefruit, red peel grapefruit and combinations thereof.
30 Particular compositions identified by way of the present invention as having significant
therapeutic value include and/or are derived from essence oil and peel oil isolated from
citrus fruit (*e.g.*, grapefruit), peel (*e.g.*, in processed form) isolated from a citrus fruit

(e.g., grapefruit), decharacterized cranberry fruit (e.g., presscake), and combinations thereof. Such compositions, and compounds derived therefrom, can then be formulated in a variety of manners, such as a dietary supplement, a pharmaceutical, or as an additive to a foodstuff. They may also include additional desirable compounds further
5 contain fats, carbohydrates, proteins, vitamins, minerals and combinations thereof.

In a related embodiment, the present invention further provides therapeutic compositions containing novel combinations and/or ratios of health-promoting compounds derived (e.g., isolated) from grapefruit and cranberry. Such compounds can be isolated from, for example, grapefruit essence oil, grapefruit peel and/or peel oil,
10 grapefruit juice, decharacterized cranberry fruit and cranberry juice. By way of illustration, such compounds can include phenolic acids, flavanoids, fibers, omega-3-fatty acids, tocochromanols, triterpenoids, ellagic acids or combinations thereof.

In a particular embodiment, the invention provides a composition containing an anthocyanin, a phenolic acid, a proanthocyanidin, or a combination thereof. In a
15 preferred embodiment, the anthocyanin content is 30% or greater of that present in the native fruit, the total phenolic acid content is 8% or greater of that present in the native fruit, and the proanthocyanidin content is 60% or greater of that present in the native fruit. Particular anthocyanins may include, for example, cyanidin-3-arabinoside, cyanidin-3-galactoside, cyanidin-3-glucoside, peonidin-3-arabinoside, peonidin-3-
20 galactoside, peonidin-3-glucoside, malvidin-3-arabinoside, malvidin-3-glucoside, or combinations thereof. Particular phenolic acids may include, for example, para-coumaric acid, caffeic acid, chlorogenic acid, ferulic acid, protocatechuic acid, cinnamic acid, benzoic acid, gallic acid, para-hydroxybenzoic acid, and combinations thereof. Particular proanthocyanidins may include, for example, flavan-3-ol polymers,
25 procyanidin B1, procyanidin B2, procyanidin B3, epicatechin oligomers, and combinations thereof. Particular flavonoids may include, for example, proanthocyanidin, flavan-3-ol, anthocyanin, flavanol, and combinations thereof. Particular flavan-3-ols may include, for example, catechin, catechin gallate, epicatechin, epicatechin gallate, epigallocatechin gallate, gallocatechin gallate, and combinations
30 thereof. Particular flavanols may include, for example, quercetin, q-3-arabinoside (avicularin), q-3-galactoside (hyperin), q-3-glucoside (isoquercitrin), q-3-rhamnoside (quercitrin), myricetin, m-3-arabinoside, m-3-rhamnoside (myricitrin), m-3-

5

10

15

20

25

30

Detailed Description of the Invention

In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

5

Definitions

The term “cancer” refers to any neoplasm (e.g., benign or malignant), such as a carcinoma (*i.e.*, usually derived from epithelial cells, *e.g.*, breast cancer) or sarcoma (usually derived from connective tissue cells, *e.g.*, a bone or muscle cancer) or a cancer
10 of the blood, such as a erythroleukemia (a red blood cell cancer) or leukemia (a white blood cell cancer). “Malignant” cancers (*i.e.*, malignancies) are generally metastatic, *i.e.*, have acquired the ability to transfer from one organ or tissue to another not directly connected, *e.g.*, through the blood stream or lymphatics.

The term “anti-cancer activity” or “anti-cancer properties” refers to the inhibition
15 (in part or in whole) or prevention of a cancer as defined herein. Anti-cancer activity includes, *e.g.*, the ability to reduce, prevent, or repair genetic damage, modulate undesired cell proliferation, modulate misregulated cell death, or modulate mechanisms of metastasis (*e.g.*, ability to migrate).

The term “hypercholesterolemia” refers to abnormally high serum levels of
20 cholesterol, typically due to defective cholesterol metabolism in a subject.

The term “anti-hypercholesterolemic activity ” refers to the ability to regulate cholesterol metabolism or reduce serum cholesterol levels in a subject.

The term “aroma” refers to the water-soluble components (*e.g.*, fraction) remaining after evaporation of a fruit juice.

25 The term “essence oil” refers to the oil-soluble components (*e.g.*, fraction) remaining after evaporation of a fruit juice.

The term “peel oil” refers to oil isolated from the peel of a citrus fruit.

The term “peel” refers to the peel of a citrus fruit which, for purposes of the present invention, may be *e.g.*, dried, shredded, or pelletized.

30 The term “citrus fruit” refers to a fruit from the genus *Citrus* that includes, *e.g.*, orange, lemon, lime, tangerine and, in particular, grapefruit (*e.g.*, pink grapefruit, red peel grapefruit).

09341-0100

5 The term "decharacterized fruit" refers to fruit from which the juice has been extracted. The decharacterized fruit can be in the form of, for example, a mash or presscake. The term "Tomah presscake" refers to a particularly preferred presscake described in U.S. Patent Nos. 5,320,861 and 5,320,861 which contains higher levels of desirable phytochemicals than are present in presscake made via conventional methods. In particular, decharacterized cranberry fruit in the form of "Tomah presscake" contains higher levels of anthocyanins, phenolic acids and proanthocyanidins than that found in presscake produced through conventional methods. For example, the anthocyanin content is typically 30% or greater of that present in native cranberry fruit, the phenolic acid content is typically 8% or greater of that present in native cranberry fruit and the proanthocyanidin content is typically 60% or greater of that present in native cranberry fruit.

The term "isolated" refers to the removal or change of a composition or compound from its natural context.

15 The term "phenolic compound" refers to a compound that is an aromatic acid having one or more hydroxyl groups on the benzene ring and naturally present at some measurable level in a fruit. In a preferred embodiment, the phenolic compounds of the invention include, *e.g.*, para-coumaric acid, caffeic acid, chlorogenic acid, ferulic acid, protocatechuic acid, cinnamic acid, benzoic acid, gallic acid, para-hydroxybenzoic acid, or a combination thereof.

20 The term "flavonoid" refers to any member of the group of aromatic, oxygen-containing, heterocyclic pigments found in the derivatives of the invention and includes for example members of the chemical subgroups 1) catechins, 2) leucoanthocyanidins and flavanones, 3) flavanins, flavones, and anthocyanins, and 4) flavonols. In preferred 25 embodiments, a flavonoid includes, *e.g.*, a proanthocyanidin, flavan-3-ol, anthocyanin, or flavanol.

The term "fiber" includes the principal chemical class of dietary fiber, which includes cellulose, hemicelluloses, gums, lignins, and preferably, pectins.

30 The term "fatty acid" refers to a fatty acid that is naturally present at some measurable level in the fruit derivatives of the invention and includes, for example, α -linolenic acid (omega-3), oleic acid (omega-9), linoleic acid (omega-6), or a combination thereof.

093514-0430
FILED IN 093514-0430

5

The term “ellagic acid” refers to a plant phenol that is naturally present at some measurable level in the fruit derivatives of the invention.

The term “foodstuff” refers to any edible substance that can be used as or in food for an animal or human. Foodstuffs include substances that may be used in the preparation of foods such as cooking oils or food additives. Foodstuffs also include dietary supplements designed to, e.g., supplement the diet of an animal or human. In addition, foodstuffs also include animals or animal products used for human consumption, such as, for example eggs or milk. Such animal themselves can be fed or treated with a composition of the invention and retain the advantageous properties of the composition (*e.g.*, decharacterized cranberry or components thereof) or impart those advantageous properties to products such as eggs or milk.

The term “pharmaceutical composition” or “therapeutic composition” refers to a composition formulated for therapeutic use.

20 The terms “health promoting”, “therapeutic” and “therapeutically active” are
used interchangeably herein, and refer to the prevention or treatment of a disease or
condition in a human or other animal, or to the maintenance of good health in a human
or other animal, resulting from the administration of a cranberry or grapefruit derivative
of the invention, or a composition derived therefrom. Such health benefits can include,
25 for example, nutritional, physiological, mental, and neurological health benefits.

Overview

The present invention is based on the identification of therapeutic cranberry and grapefruit products (e.g., derivatives), and compounds isolated from the products, 30 having novel therapeutic and/or health promoting values (see Tables 1 and 2). In particular, therapeutic cranberry and grapefruit products of the invention are shown

herein to exhibit significant anti-cancer and anti-hypercholesterolemia activity when administered to a subject *in vivo* and when tested *in vitro*.

In a particular embodiment, therapeutic methods of the invention employ decharacterized cranberry fruit, preferably Tomah presscake, which is enriched for a number of health promoting compounds and exhibits e.g., significant anti-cancer properties when administered to a subject. Moreover, a variety of particular health promoting compounds derived from decharacterized cranberry have been identified and are discussed below.

In another embodiment, therapeutic methods of the invention employ grapefruit products (derivatives) which are novel sources of compounds having significant therapeutic value, in, for example, the prevention or treatment of cancer by, e.g., lowering rates of metastasis. In addition, as described herein, a subset of these grapefruit derivatives are also enriched with compounds suitable for treating disease related to high cholesterol (hypercholesterolemia).

In another embodiment, therapeutic methods of the invention employ compounds derived from cranberry and grapefruit juices which, as shown herein, have anticancer activity (e.g., reduced metastasis rates) e.g., when prepared in concentrated form and administered to a mammal *in vivo*.

Accordingly, the identification of particular beneficial compounds in cranberry and grapefruit derivatives has allowed for the development of convenient methods and compositions (e.g., formulations) for administering therapeutic compounds to treat or prevent particular diseases. Moreover, the therapeutic compounds and compositions described herein have the additional advantage of being readily manufactured into palatable forms (e.g., as foodstuffs such as juices and food bars or as dietary supplements) for convenient oral administration.

Methods for obtaining and preparing cranberry and grapefruit products of the invention, identifying (e.g., characterizing) and obtaining therapeutic components of the products, evaluating biological activity *in vitro* and *in vivo* of the products and components, and methods of using the products and novel compositions containing the products or combinations of components isolated from the products, are discussed in the appropriate subsections below.

Table 1. Summary Table of Effectiveness of Different Cranberry Derivatives

	Sample	anti-cancer	anti-hypercholesterolemic	type of study
1	cranberry essence (OS2)	-	-	cells
2	decharacterized cranberry	+	-	animals
3	cranberry juice, 2x	+	-	animals

5

Table 2. Summary of Effectiveness of Different Grapefruit Derivatives

	Sample	anti-cancer	anti-hypercholesterolemic	type of study
4	grapefruit aroma (OS1)	-	-	cells
5	grapefruit essence oil (OS3)	+	+	cells
6	grapefruit red peel oil (OS4)	+	+	cells
7	grapefruit peel processed	-	+	animals
8	grapefruit juice, pink, 2x	+	-	animals

DECLASSIFIED

Methods for Preparing Cranberry and Grapefruit Derivatives

Cranberry and grapefruit products (i.e., derivatives) of the invention may be isolated from whole cranberry, grapefruits, or juices, peels, rinds thereof, using any suitable art recognized method. Preferred derivatives include cranberry essence, cranberry juice (e.g., concentrated), decharacterized cranberry fruit, grapefruit essence oil, grapefruit peel oil, grapefruit peel (e.g., processed), and grapefruit juice (e.g., concentrated).

In a particular embodiment, decharacterized cranberry fruit is obtained using the method described in U.S. Patent Nos. 5,320,861 and 5,419,251, hereby incorporated by reference.

In all cases, the cranberry or grapefruit derivatives are preferably obtained in a form suitable for use in a foodstuff, dietary supplement, or pharmaceutical composition. Further, it is understood that with regard to any of the techniques for preparing a cranberry or grapefruit derivative described herein, it may also be desirable to avoid exposing the derivative, or component thereof, to oxygen by, e.g., protective blanketing of the derivative or component with carbon dioxide or nitrogen gas, or by, e.g., exposing the derivative or component, where appropriate, to BHT, ascorbic acid, low temperature, or a combination of these conditions.

Cranberry and Grapefruit Derivatives

As part of the present invention, several cranberry derivatives, including cranberry essence (evaporated juice), concentrated cranberry juice, and decharacterized cranberry fruit were analyzed for health promoting compounds (see Table 2). These cranberry derivatives were analyzed using both chemical analysis and bioactivity assays as described herein. In addition, a number of grapefruit derivatives, including grapefruit aroma, grapefruit essence oil, grapefruit red peel oil, processed grapefruit peel, and grapefruit juice also were studied for their *in vitro* and *in vivo* therapeutic activity and analyzed for health promoting compounds (see Table 1).

Accordingly, by way of the studies described herein, it was shown that particular cranberry and grapefruit are novel sources of therapeutically beneficial compounds such as a phenolic acid, flavanoid, pectin, omega-3-fatty acid,

tocochromanol, triterpenoid, ellagic (see also Table 15). Decharacterized cranberry, particularly "Tomah presscake", provides several advantages over currently known sources of such therapeutically beneficial compounds including, for example, a remarkably high concentration of particularly desirable components (*e.g.*, an anthocyanin, a phenolic acid, a proanthocyanidin). Accordingly, the cranberry and grapefruit derivatives of the invention, or components thereof, can be used in foodstuffs, or as dietary supplements or pharmaceutical compositions.

Thus, in one embodiment, the invention provides a cranberry or grapefruit derivative, or a composition comprising one or more components isolated from the cranberry or grapefruit derivative, *e.g.*, see Tables 12-14, which promotes health in a human or other animal. The cranberry or grapefruit products of the invention, and compositions derived therefrom, can additionally contain one or more exogenous (*i.e.*, externally added) compounds to further enhance the therapeutic value of the derivative or composition derived therefrom, for example, by acting in synergism with one or more native components of the derivative. For example, grapefruit derivatives of the invention (and compounds isolated therefrom, such as a flavanone, flavone, liminoid, fiber, essential oil, or glucaric acid) having bioactivity may be used alone or in combination with cranberry derivatives of the invention (such as decharacterized cranberry fruit and compounds isolated therefrom) for the prevention or treatment of a disease or condition in a human or other animal, or for the maintenance of good health in a human or other animal. Such health benefits can include, for example, nutritional, physiological, mental and neurological health benefits.

Phenolic Compounds

In particular, the cranberry and grapefruit derivatives of the invention contain one or more phenolic compounds, such as those listed in Table 13.

Such phenolic compounds can act as potent antioxidants and, therefore, can prevent or delay oxidation reactions which cause various diseases. Accordingly, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom can be used as anti-oxidants. For example, they can inhibit lipid peroxidation, scavenge free radicals and active oxygen, inactivate lipoxxygenase, and chelate iron ions. They also can be used to inhibit erythrocyte aggregation and

sedimentation. Moreover, epidemiological studies have demonstrated that the consumption of phenolic compounds is associated with a reduced risk of cancer.

Accordingly, the cranberry and grapefruit derivatives of the invention and components derived therefrom (*e.g.*, fractions rich in phenolic compounds) can be used
5 to treat cancer or cholesterol-related disorders with fewer side effects compared to standard chemotherapies.

Flavonoids

Cranberry and grapefruit derivatives of the invention also contain one or more
10 flavonoids, such as those listed in Table 14.

Flavonoids have widespread anti-cancer and anti-hypercholesterolemic properties. For example, a flavonoid, in particular, a proanthocyanidin extract, has been found to inhibit a key enzyme associated with cell proliferation and skin cancer (Bosmer
15 *et al.*, Planta Med 62:212-6 (1996)). In addition, the flavonoid quercetin can inhibit tumor production by chemical carcinogens, inactivate some enzymes involved in the metabolism of carcinogens, and inhibit LDL oxidation.

Accordingly, the cranberry and grapefruit derivatives of the invention and components derived therefrom (*e.g.*, fractions rich in flavonoid compounds) can be used
20 to treat cancer and cholesterol-related disorders.

Fiber

Cranberry and grapefruit derivatives of the invention also contain fiber, in particular, pectin, and fiber when administered to animals or humans increases the bulk of the feces and this can relieve, *e.g.*, constipation and may reduce the incidence of
25 diverticula forming in the large intestine of older people. Moreover, there is some evidence that a diet high in fiber can reduce colon cancer and reduce plasma cholesterol concentration.

Accordingly, the cranberry and grapefruit derivatives of the invention and components derived therefrom (*e.g.*, fractions rich in pectin) can be used to treat the
30 above-mentioned disorders.

Fatty Acids

Cranberry and grapefruit derivatives of the invention also contain beneficial fatty acids. Fatty acids, for example, omega-3 fatty acids, are essential for growth and development throughout the life cycle. For example, omega-3 fatty acids are known to play an important role in, 1) the normal function of the retina and brain, especially in new born infants, 2) maintaining favorable serum triglycerides in normal subjects and in patients with hypertriglyceridemia, 3) the normal function of the vascular and neurological systems, and 4) reducing LDL (low density lipoprotein) cholesterol in patients with hyperlipidemia (provided that the saturated fatty acid content in the diet is decreased).

Beneficial fatty acids, also derivable from the derivatives of the invention, play important roles in normal physiological functions including, *e.g.*, overall growth, healthy skin, reproduction, and cardiovascular health.

Accordingly, formulations can be prepared by those of ordinary skill in the art containing a desirable fatty acid derived from a cranberry or grapefruit derivative. Such formulations have application in the medical and pharmaceutical industries for enhancing, maintaining or treating any of the above-mentioned biological functions or disfunctions. In addition, given the wide spectrum of biologic processes affected by these fatty acids, the derivatives of the invention can also be used as a food additive or dietary supplement.

For example, in the food industry, to raise the availability of desirable fatty acids in a consumer's diet, the derivatives of the invention, or compositions derived therefrom (*e.g.*, containing components or fractions thereof), can be added to, for example, juices, bakery products, infant formulas, *etc.* As dietary supplements, the derivatives of the invention or compositions derived therefrom can be taken in the form of *e.g.*, liquids, pills, or capsules as are known in the art. As discussed further below, methods for formulating such vehicles of administration can be performed using standard techniques.

In another embodiment, the derivatives of the invention or compositions derived therefrom (*e.g.*, containing health-promoting fatty acids) can be fed or otherwise administered to laying hens to produce eggs rich in desirable fatty acids, or to cows or other livestock to produce meat and dairy products rich in such fatty acids. The

resultant food products derived from these animals can then be consumed by humans for their enhanced nutritional and health benefits.

Alternatively, the derivatives of the invention or compositions derived therefrom can be fed or otherwise administered to animals, such as pets or domesticated livestock, for therapeutic purposes (*e.g.*, to correct problems such as dry skin, allergic reactions, and cancer).

Tocochromanols

Cranberry derivatives of the invention, in particular, also contain a remarkably high concentration of tocochromanols (a class of compounds that includes tocopherols and tocotrienols), such as α -tocopherol, γ -tocopherol, δ -tocopherol, α -tocotrienol, γ -tocotrienol, δ -tocotrienol, or a combination thereof. A large body of research has shown the importance of tocopherols and tocotrienols in the defense against numerous biological disorders, for example, breast cancer, as described in 60/137,405 by Nawar, W., hereby incorporated by reference.

Tocochromanols are one of the active agents in vitamin E which has been recognized as an anticarcinogenic agent for a number of years (Haenszel *et al.*, *Int. J. Cancer*, 36:43-48 (1985); Menkes *et al.*, *N. Engl. J. Med.*, 315:1250-1204 (1986); Stahelin *et al.*, *Ann. NY Acad. Sci.*, 570:391-399 (1989)). In addition, *in vitro* and *in vivo* studies, including human studies, have demonstrated that vitamin E interferes with the development of carcinogenesis that results from exposure to various environmental factors known to enhance oxidant stress (Borek *et al.*, In, *Mechanisms of cellular transformation by carcinogenic agents*, New York, Pergamon (1987), Borek *et al.*, In, *Medical, biochemical and chemical aspects of free radicals*, Amsterdam, Elsevier, (1989); Borek *et al.*, *Proc. Natl. Acad. Sci. USA* 83:1490-1494 (1986); *Proc. Natl. Acad. Sci. USA*, 88:1953-1957 (1991)). In addition, α -tocopherol, a component of vitamin E, is a hydrophobic, peroxy radical trapping, chain-breaking antioxidant found in biological membranes. Accordingly, the protective role vitamin E plays in inhibiting a variety of human malignancies is mainly attributed to its components having the ability to protect the lipid material of the organs against oxidation (Ames *et al.*, *Science* 230:271-279 (1987); Doll *et al.*, *J. Natl. Cancer Inst.* 66:1193-1194 (1981); Greenwald

et al., *Cancer* 65:1483-1490 (1990); Menzel *et al.*, *J. Agr. Food Chem.*, 20:481-486 (1972)).

Accordingly, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom (*e.g.*, fractions rich in tocochromanols) can be used to
5 treat respiratory, inflammatory, neurological, dermatological, ophthalmological, and gastroenterological diseases.

Triterpenoids

In addition, cranberry and grapefruit derivatives of the invention contain one or more triterpenoids, such as, *e.g.*, ursolic acid. Such triterpenoids are known to confer significant health benefits, *e.g.*, as antiinflammatories or hepatoprotectants. For example, ursolic acid is effective in protecting against chemically induced liver injury in laboratory animals and also has antihyperlipidemic properties. Ursolic acid has also been noted for its antitumor-promotion effects (Liu, J Ethnopharmacol 49:57-68 (1995)).

Accordingly, cranberry or grapefruit derivatives of the invention, and components derived therefrom (*e.g.*, fractions rich in triterpenoids), can be used to, *e.g.*, inhibit inflammation, treat liver disorders, and inhibit tumor promotion.

20 Ellagic Acid

In addition, cranberry and grapefruit derivatives of the invention also contain a plant phenol such as ellagic acid. Plant phenols such as ellagic acid have anticancer properties and ellagic acid in particular, can protect animals against benzo[a]pyrene-induced neoplasia (Lesca, P., Carcinogenesis 4:1651-3(1983)).

25 Accordingly, cranberry or grapefruit derivatives of the invention and components derived therefrom (*e.g.*, fractions rich in ellagic acid) can be used as an anti-cancer agent.

Methods for Isolating, Identifying, and Analyzing Specific Components from
Cranberry and Grapefruit Derivatives

To isolate and analyze constituent therapeutic components (compounds) from cranberry and grapefruit derivatives, a variety of art-recognized techniques and assays

can be employed. For example, as described in the studies provided herein, oil-based samples (*e.g.*, grapefruit essence oil or grapefruit red peel oil) can be prepared for analysis by converting the fatty acids in the oil to their methyl esters, for example, by refluxing with MeOH/MeO⁻ Na⁺. The resultant methyl esters can then be analyzed, *e.g.*,
5 by gas chromatography.

Phenolic compounds of the cranberry and grapefruit derivatives of the invention can be analyzed and extracted using HPLC analysis and solvent extraction, respectively. The isolated extracts can be dissolved in hexane and then extracted with a methanol/water solution followed by centrifugation. The extract can then be dried, and
10 the residue can be resuspended in methanol/water for HPLC analysis.

For example, tocochromanols contained in an oil fraction derived from a cranberry or grapefruit derivative of the invention can be separated and analyzed using, for example, the methods of Carpenter (Carpenter, Jr., A.P., *J. Amer. Oil Chemists' Soc.*, 56:668 (1979)).

15 Triterpenoids can be extracted and analyzed using, for example, thin layer chromatography and high-performance liquid chromatography. For example, an the isolated oil fraction derived from a cranberry or grapefruit derivative can be saponified with KOH, the unsaponifiables extracted with ether, and the resultant material can be fractionated on thin-layer chromatography (TLC) plates where the individual bands that
20 are subsequently resolved can be scraped and extracted with a chloroform/methanol solvent. These resultant samples can then be analyzed using, *e.g.*, gas and high-performance liquid chromatography (HPLC).

Other methods known in the art may also be employed, in place of or in combination with, the methods described above for isolating cranberry or grapefruit
25 components, particularly to "scale up" the quantity of the isolated components. For example, chromatographic techniques may be used for isolating components of the cranberry or grapefruit derivatives of the invention, in sufficient and pure quantities, such that the component may be administered alone or as part of a composition or product described herein (*e.g.*, foodstuffs, dietary supplements, pharmaceuticals, *etc.*).
30 In particular, gas liquid chromatography, gas solid chromatography, high pressure or high performance liquid chromatography (HPLC) (*e.g.*, normal, reverse, or chiral), ion exchange chromatography, or size exclusion chromatography can be employed as

described, for example, in *Advances in Chromatography*, Brown, Eds., Marcel Dekker, Pub. (1998); *Basic Gas Chromatography*, Harold *et al.*, John Wiley & Sons, Pub. (1997); *Column Handbook for Size Exclusion Chromatography*, Wu, Ed., Academic Press, Pub. (1999); *Fundamentals of Preparative and Nonlinear Chromatography*, Guichon *et al.*, Eds., Academic Press, Pub. (1994); *Handbook of Process Chromatography: A Guide to Optimization, Scale-Up and Validation*, Hagel *et al.*, Eds., Academic Press, Pub. (1997); *HPLC Methods for Pharmaceutical Analysis*, Lunn *et al.*, John Wiley & Sons, Pub. (1997); and *Practical High-Performance Liquid Chromatography*, Meyer, Wiley-Liss, Pub. (1999), each of which is incorporated by reference herein. Such isolated components, which can be separated as “value added” fractions (*e.g.*, fractions having therapeutic value), are typically rich in at least one beneficial component identified from a cranberry or grapefruit derivative described herein. These isolated components or fractions may be further combined to provide a composition rich in more than one component or, *e.g.*, a desired combinations thereof.

15 In addition, a particular formulation intended for the treatment or prevention of a particular disease or condition may be formulated to be rich in those components having a therapeutic effect on the disease or condition (*e.g.*, associated with affecting a change in any of the mechanisms associated with that particular disease or condition). For example, a formulation suitable for administering to a subject with cancer is preferably

20 rich in cranberry or grapefruit derived components having antioxidant and other anti-cancer properties, whereas a formulation for administering to a subject with a dietary need, may be rich in, for example, beneficial fatty acids.

In Vitro Methods for Evaluating the Therapeutic Properties of Cranberry and Grapefruit Derivatives And Components Derived Therefrom

25 The health promoting properties, *e.g.*, anticancer activity, of cranberry and/or grapefruit derivatives of the invention, and compositions derived therefrom, can be evaluated using a variety of art-recognized cell-based assays, *e.g.*, cell proliferation assays using tumor cells.

30 For example, in one embodiment, a tumor cell proliferation assay is performed by measuring the incorporation of [³H] thymidine into the DNA of dividing cells, as is known in the art. For example, a solution containing cranberry and/or grapefruit

derivatives of the invention, or components derived therefrom (*e.g.*, a phenolic acid rich fraction rich fraction), can be added to tissue culture plates, for example, in decreasing concentrations and incubated at 37°C for 3 days, after which tritiated thymidine is added to each well to determine the number of dividing cells at each concentration. The cells
5 are further incubated for a sufficient period of time, *e.g.*, 4 hrs, to allow for the incorporation of a detectable radiolabel into the DNA of dividing cells and then medium and excess label are removed. The cells can then be harvested by, *e.g.*, trypsinization, and the amount of radioactivity present in the cells is measured using standard techniques. The concentration at which the cranberry or grapefruit derivatives of the
10 invention exhibit 50% or 90% inhibition of cell growth (respectively, IC50 or IC90) is determined by comparing the radioactivity measured in the extract-treated cells as compared to untreated control cells.

Another method for determining the viability of tumor cells after exposure to an appropriate cranberry and/or grapefruit derivative of the invention, or a component
15 derived therefrom, employs a vital dye (3-[4-5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT)) that, when mixed into a cell sample, exhibits a detectable signal that distinguishes viable from non-viable cells. The intensity of the blue color, due to a formazan product formed by cellular reduction of MTT by the mitochondrial dehydrogenase of the surviving cells, is then measured as an indication of the viability
20 of the remaining cells (Hansen *et al.*, *J. Immunol. Methods*, 119:203-210 (1989)). Percent viability can be determined by relating absorbance/concentration of the treated cells to that of the non-treated controls.

To resolve the long term growth effects on cells caused by exposure to an appropriate cranberry and/or grapefruit derivative of the invention, or a component
25 derived therefrom, can be determined by incubating plates containing the cell culture medium plus the MTT reagent at its IC50 concentration at 37°C. Plates are removed at appropriate intervals, the medium aspirated, the cells trypsinized, resuspended, counted with a hemocytometer, and the number of cells plotted against time to construct growth curves.

In Vivo Methods for Evaluating the Therapeutic Properties of Cranberry and Grapefruit Derivatives And Components Derived Therefrom

In another embodiment, cranberry and/or grapefruit derivatives of the invention, and compositions derived therefrom, can be tested for their *in vivo* therapeutic effect by administering (e.g., orally) the extracts or compositions in a suitable form (e.g., as a food stuff, dietary supplement, or pharmaceutical composition) to a human or other animal, and then observing the physiological effect (e.g., compared to a control). The human or animal can be, for example, suffering from a disease or condition, such as those described herein (e.g., cancer or hypercholesterolemia). Thus, a reduction in the physical symptoms of the disease can be measured as an indication of the therapeutic efficacy of the cranberry and/or grapefruit derivatives or compositions derived therefrom.

In another approach for evaluating anti-tumor activity, cranberry and/or grapefruit derivatives of the invention or compositions derived therefrom (e.g., a fraction thereof) can be used in a controlled animal study where tumors are induced in the animal via diet, by applying a chemical tumor promoter to the skin, or by the implantation of tumor cells in the presence or absence of the test agent (see, e.g., Example 2). Various assays, such as those described below, can then be used to examine the progression of carcinogenesis in the presence or absence of the administration of the extracts or compositions of the invention.

Methods of Use

Treatment of Cancer

In one embodiment, cranberry and/or grapefruit derivatives of the invention, and compositions derived therefrom (particularly those containing detectable amounts of, e.g., anthocyanins, phenolics, proanthocyanidins, tocotrienols, or flavonoids), are administered to a mammal (e.g., a human or animal) to treat or prevent cancer. Such derivatives and compositions also can be administered in combination with other anti-cancer agents. In particular, the cranberry and grapefruit derivatives of the invention, and compositions derived therefrom, can be administered separately or together and/or with either tamoxifen and/or a flavonoid for the treatment of, for example, breast cancer. These combinations of agents encompassed by the invention are particularly effective

because of the ability of tocochromanols to act in synergy with tamoxifen and/or flavonoids in the inhibition of tumorigenic cells.

For example, it is known that most breast cancers consist of hormone-dependent as well as hormone independent cells. The drug tamoxifen, a synthetic non-steroidal estrogen antagonist, has been widely used in the treatment of hormone-responsive breast cancer. In addition, the inhibitory effects of various combinations of the palm oil tocotrienol-rich fractions as well as individual tocotrienols in combination with tamoxifen on at least two breast cancer cell lines (*i.e.*, estrogen receptor-negative MDA-MB-435 and estrogen receptor-positive MCF-7) have been demonstrated (Guthrie, *et al.*, *Asia Pacific J. Clin. Nutr.* 41-45 (1997)).

Prior to the present invention, treatment of cancer patients with tamoxifen had several drawbacks. For example, tumors can develop resistance to tamoxifen, possibly caused by the drug's intrinsic estrogen antagonist properties (Osborne *et al.*, *J. Natl. Cancer Inst.* 87:746-750 (1995)). Also, tamoxifen may increase the incidence of new primary malignancies, *e.g.* endometrial, liver, and colorectal cancers (Rutgrist *et al.*, 1995). Accordingly, the present invention provides the advantage of enabling the administration of tamoxifen in lower doses, for example, in combination with a cranberry derivative of the invention or a composition derived therefrom (particularly one having a high tocotrienol content) to avoid these undesirable effects.

Treatment of High Cholesterol and Diseases Arising Therefrom

In another embodiment, the cranberry and grapefruit derivatives of the invention, and compositions derived therefrom (*e.g.*, those having high tocochromanol and/or phytochemical content), can be used to treat or prevent high cholesterol and related diseases such as arteriosclerosis and heart disease. Indeed, the efficacy of various tocochromanol and plant derived phytochemicals in reducing cholesterol levels in animals, including humans, is well supported in the scientific literature.

Animal studies have shown that tocotrienol-containing bran oil, barley oil, and palm oil suppressed cholesterologenesis when fed to chicken. In addition, hypercholesterolemic pigs fed the TRF-supplemented diet showed a 44% decrease in total serum cholesterol and a 60% decrease in LDL cholesterol, with the decrease

persisting for 8 weeks even after putting the animals back on the control corn-based diet (Quereshi *et al.*, In, *International Palm Oil Conference*, pp 45-47 (1988)).

Similarly, a study involving 47 hypercholesterolemic subjects administered dietary supplements containing 200 mg of TRF per day for 4 weeks resulted in, respectively, a 15-22% and 10-20% reduction in serum total and LDL cholesterol (Quereshi, *et al.*, *Lipids* 20:817-824 (1985); Quereshi, *et al.*, *Am. J. Clin. Nutr.*, 53:1021S-1026S (1991)). In addition, in studies where the hypocholesterolemic effect of tocotrienols was compared with that of other drugs, the tocotrienols were more effective. For example, in a study involving chickens, T3 was demonstrated as being twice as effective as LovastatinTM, a drug currently used for cholesterol control in humans. And most of the drugs most commonly used today in the therapy of hypercholesterolemia (*i.e.*, Nicotinic acid (Grundy *et al.*, *J. Lipid Res.*, 22:24-36 (1981)), CompactinTM and LovastatinTM (Illingworth *et al.*, *Eur. Heart J.*, Supp. E:103-111 (1987); Endo *et al.*, *Biotechnology*, 26:301-320 (1994)), CholestyramineTM and ColestipolTM (Shepherd *et al.*, *Biochem. Soc. Trans.* 15:199-201 (1980)), ClofibrateTM and GemfibrosilTM (Kesaniemi *et al.*, *JAMA*, 251:2241-2246 (1984)) and ProbucolTM are known to produce various side effects (Illingworth *et al.*, *Am. J. Cardiol.*, 60:33G-42G (1987)). In contrast, no toxic effects were observed in the studies where tocotrienols were administered.

Accordingly, the cranberry and grapefruit derivatives of the invention, and compositions derived therefrom, can be used in the treatment of high cholesterol (cholesterolemia) and other associated conditions such as heart disease.

Treatment of Other Diseases and Disorders

In yet another embodiment, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom (particularly those having high, *e.g.*, a high tocochromanol content) can be used in the treatment or prevention of a wide range of other diseases and disorders that include aging, respiratory, inflammatory, neurological, dermatological, ophthalmological, and gastroenterological diseases. Indeed, a large volume of reported research provides evidence that vitamin E-containing tocochromanols plays a critical role in the above-mentioned conditions. In addition, a number of studies indicate that citrus-derived flavanones, flavones, liminoids, fiber (*e.g.*,

pectin), essential oils, and glucaric acid have, *e.g.*, anticancer-activity and/or anti-hypercholesterolemic activity (for a review, see Johns *et al.*, *Recent Advances in Phytochemistry*, pp.31-52, Plenum Press (1997).

For example, cranberry or grapefruit derivatives of the invention and
5 compositions derived therefrom can be used to prevent endothelial injury, such as ischemic and reperfused myocardium and ulcers. In addition, the extracts and compositions can be used to inhibit tumor necrosis factor biosynthesis that, in turn, decreases inflammation (*e.g.*, by inhibiting respiratory bursts of neutrophils or via free radical scavenging). Accordingly, cranberry or grapefruit derivatives of the invention
10 and compositions derived therefrom (particularly those having high tocochromanol and/or flavanoid content) can be used as anti-inflammatory agents for the prevention and treatment of a wide variety of diseases and conditions involving minor, acute and chronic inflammation.

Cranberry or grapefruit derivatives of the invention and compositions derived
15 therefrom (particularly those having high tocochromanol (particularly those having high tocochromanol and/or flavanoid content) also can be used to treat glucose intolerance in diabetes mellitus, and/or to restore acute glucose-induced insulin response in non-insulin-dependent diabetes mellitus.

In addition to the above-stated uses, cranberry or grapefruit derivatives of the
20 invention and compositions derived therefrom (particularly those having high tocochromanol and/or flavanoid content) can be used to enhance the immune response in animals and humans, for example, by reducing the amount of fatty acids in biological tissues. Since fatty acid levels effect the immune system, the compounds of this invention may serve as immunoregulators. They may, for example, be used to increase
25 antibody titers to foreign proteins.

In addition, the reduction in fatty acid, cholesterol, fatty acid and/or glucose levels induced by the compounds of the invention can be obtained without attendant substantial weight loss, resulting in an increased feed to protein conversion ratio. Therefore, the extracts and compositions of the invention can be used to increase feed
30 conversion efficiency.

098341-0430
102410-225860

Thrombotic diseases and conditions that may be treated using cranberry or grapefruit derivatives of the invention and compositions derived therefrom include, but are not limited to, pulmonary disease (for example, involving reduced conductance, compliance, or constriction), excessive fluid accumulation or pulmonary edema, respiratory distress, asthma, pulmonary vascular permeability, pulmonary vasoconstriction, pulmonary hypertension, pulmonary embolism, cardiac ischemia, myocardial infarction, cardiopulmonary bypass associated dysfunction, vasoconstriction, organ dysfunction, platelet dysfunction, cardiac disease, chronic obstructive arterial disease caused by arteriosclerosis, vasoconstriction, renal artery stenosis, myocardial infarction, stroke, deep vein thrombosis, peripheral arterial occlusion, and other blood system thromboses.

The antioxidizing properties of the cranberry or grapefruit derivatives of the invention and compositions derived therefrom may also be applied to, but are not limited to, the treating and preventing of cancerous conditions by, for example, preventing or limiting cancer-causing mutations in the genetic material of an animal or a human.

Antiatherogenic diseases and conditions that can be treated using cranberry or grapefruit derivatives of the invention and compositions derived therefrom include, but are not limited to, atherosclerosis, arteriosclerosis, myocardial infarction, ischemia (*i.e.*, myocardial ischemia, brain ischemia, and renal ischemia) and strokes.

Inflammatory diseases and conditions that can be treated using cranberry or grapefruit derivatives of the invention and compositions derived therefrom include, but are not limited to, essential hypertension, hypertension of congestive heart failure, renal dysfunction caused by reduced myocardia output, endotoxemia, chronic liver disease or hypertension, pulmonary inflammation in asthma, lung injury (bronchitis, pneumonia, or acute); rheumatic diseases (for example, rheumatoid arthritis or systemic lupus erythematosus), inflammatory bowel disease (for example, ulcerative colitis), irritable bowel disease (such as villous adenoma), gastrointestinal disorders caused by excess

acids, pepsin or bile salts, Zollinger-Ellison syndrome, skin diseases or trauma (such as burns or acid or caustic injury), gout, Bartter's syndrome, fever, rheumatoid diseases, pain, and functio laesa.

Immunoregulatory diseases and diseases that can be treated using cranberry or
5 grapefruit derivatives of the invention and compositions derived therefrom include, but are not limited to, autoimmune diseases, for example, AIDS, chronic fatigue syndrome, graft rejections, and other viral diseases that impair the immune system.

Synergy with Other Components Derived From Cranberry or Grapefruit Derivatives
10 and/or Exogenous Compounds

In another embodiment, cranberry and/or grapefruit derivatives of the invention, or one or a combination of components derived therefrom, are administered to a subject with an additional (exogenous) compound, *e.g.*, an anti-cancer such as tamoxifen, and/or in combination with a flavonoid for the treatment or prevention of cancer. These
15 combinations of agents encompassed by the invention are particularly effective because of their known ability to act in synergy in the inhibition of tumorigenic cells. The flavonoid may be contained in or derived from cranberry or citrus fruit such as, *e.g.*, a grapefruit.

Flavonoids are polyphenolic compounds which occur in plant foods, particularly
20 citrus. These compounds include the flavones, *e.g.* tangeretin; the flavanones, *e.g.* hesperetin; the isoflavones, *e.g.* genistein; and the flavonols, *e.g.* quercetin and these are predicted to be present at various levels in one or more of the grapefruit derivatives of the invention. It is understood that with no more than routine experimentation in combination with the teachings of the specification the skilled artisan would be able to
25 determine which cranberry or grapefruit derivatives of the invention have desirable levels of a polyphenolic compounds suitable for administering to a mammal alone or in combination with another derivative of the invention or exogenous compound. Several studies have demonstrated the anticancer properties of flavonoids from various plant sources (Cook *et al.*, *J. Nutr. Biochem.* 7:66-76 (1996); Hertog *et al.*, *Nutr. Cancer*
30 20:21-29 (1993); Middleton *et al.*, *Trends Pharm. Sci.*, 5:335-338 (1984)). Further, various combinations of flavonoids from different sources have been shown to be

synergistic in their ability to inhibit the proliferation of a breast cancer cell line (MDA-MB-435 cells).

In particular, synergistic effects between the tocochromanols and flavonoids, with γ -T₃ and tangeretin being the most effective combination, have been observed
5 when tested for their ability to inhibit growth in MDA-MB-435 and MCF-7 breast cancer cells (IC₅₀ 0.05 μ g/mL and 0.02 μ g/mL, respectively) (Guthrie *et al.*, *Asia Pacific J. Clin. Nutr.* 6:41-45 (1997)). In addition, with few exceptions, combinations (1:1:1) of tocotrienols, flavonoids, and tamoxifen were more effective than 1:1 combinations of T₃ and flavonoids, T₃ and tamoxifen, or flavonoids, and tamoxifen.
10 The most potent combinations are γ -T₃/tangeretin/tamoxifen with the MDA-MB-435 cells and δ -T₃ /hesperetin/tamoxifen with the MCF-7 cells.

Accordingly, cranberry and grapefruit derivatives of the invention and compositions derived therefrom (particularly those having high tocochromanol and/or flavonoid content) can be used alone or in combination with tamoxifen and/or
15 flavonoids as potent anti-cancer agents.

Formulations and Methods of Administration

The cranberry and grapefruit derivatives of the invention and components derived therefrom can be administered to a subject in any suitable form. For example,
20 the extracts and compositions of the invention are sufficiently stable such that they can be readily prepared in a form suitable for adding to various foodstuffs including, for example, juice, fruit drinks, carbonated beverages, breakfast cereals, biscuits, cakes, muffins, cookies, toppings, bread, bagels, fiber bars, soups, crackers, baby formulae, salad dressings, cooking oils, and meat extenders.

25 In addition, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom can be formulated as a pharmaceutical composition (e.g., a medicinal drug) for the treatment of specific disorders.

In another embodiment, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom can be formulated as a dietary supplement.

30 Suitable additives, carriers, and methods for preparing such formulations are well known in the art.

00344-0444

For example, pharmaceutical compositions may take the form of tablets, capsules, emulsions, suspensions and powders for oral administration, sterile solutions or emulsions for parenteral administration, sterile solutions for intravenous administration and gels, lotions and cremes for topical application. The pharmaceutical compositions may be administered to humans and animals in a safe and pharmaceutically effective amount to elicit any of the desired results indicated for the compounds and mixtures described herein. In addition, the extracts of the invention may be used in cosmetics.

The pharmaceutical compositions of this invention typically comprise a pharmaceutically effective amount of a cranberry and/or grapefruit derivative, or fraction thereof, containing, for example, a phenolic acid, flavonoid, pectin, omega-3-fatty acid, tocochromanol, triterpenoid, ellagic acid, or combination thereof (as pertains to cranberry, see also Table 15), and if suitable a pharmaceutically acceptable carrier. Such carriers may be solid or liquid, such as, for example, cornstarch, lactose, sucrose, olive oil, or sesame oil. If a solid carrier is used, the dosage forms may be tablets, capsules or lozenges. Liquid dosage forms include soft gelatin capsules, syrup or liquid suspension.

Therapeutic and prophylactic methods of this invention comprise the step of treating patients or animals in a pharmaceutically acceptable manner with the compositions and mixtures described herein. As used herein, the term "therapeutically effective amount" refers to an amount effective to achieve a desired therapeutic effect, such as, achieving desirable levels of anti-cancer activity (*e.g.*, reduction of tumor size, incidence of metastasis) or anti-cholesterol activity (*e.g.*, lower blood levels of LDL-cholesterol and total serum cholesterol and higher ratios of HDL-cholesterol to LDL-cholesterol, reduced apoB secretion). The pharmaceutical compositions of this invention may be employed in a conventional manner for the treatment and prevention of any of the aforementioned diseases and conditions. Such methods of treatment and prophylaxis are well-recognized in the art and may be chosen by those of ordinary skill in the art from the available methods and techniques. Generally, dosage ranges may be from about 1 to about 1000 mg/day. However, lower or higher dosages may be employed. The specific dosage and treatment regimens selected will depend upon

factors such as the patient's or animal's health, and the severity and course of the patient's (or animal's) condition and the judgment of the treating physician.

The cranberry and grapefruit derivatives of the invention and compositions derived therefrom also can be used in combination with conventional therapeutics used
5 in the treatment or prophylaxis of any of the aforementioned diseases. Such combination therapies advantageously utilize lower dosages of those conventional therapeutics, thus avoiding possible toxicity incurred when those agents are used alone.

In foodstuffs, the cranberry and grapefruit derivatives of the invention and compositions derived therefrom can be used with any suitable carrier or edible additive.
10 For example, the cranberry and grapefruit derivatives of the invention may be used in a variety of foodstuffs, such as drinks, for example, juice drinks, sports drinks, and drink mixes. Advantageously, the above-mentioned foodstuffs may be included in low fat, low cholesterol, or otherwise restricted dietary regimens.

Pharmaceutical compositions, dietary supplements, and foodstuffs of the present
15 invention can be administered to humans and animals such as, for example, livestock and poultry. Once an animal has consumed or otherwise been administered the composition, it can advantageously retain the hypocholesterolemic, anti-cancer, or other advantageous biological activities of the administered compounds. Accordingly, an animal raised under these conditions, or any product derived therefrom, such as, for
20 example, milk, may be consumed by a human or another animal to derive the benefits of the derivatives of the invention or compositions derived therefrom. For example, a chicken which ingests feed fortified with the derivatives of the invention may later be eaten by a human to derive the cholesterol-reducing benefits.

25 This invention is further illustrated by the following examples which should not be construed as limiting.

EXAMPLE 1***IN VITRO* DEMONSTRATION OF THE ANTI-CANCER PROPERTIES OF CRANBERRY AND GRAPEFRUIT DERIVATIVES**

The following studies were performed to examine the anti-cancer properties of
 5 cranberry and grapefruit derivatives.

Four different cranberry and grapefruit extracts were prepared using methods described herein and tested for their ability to inhibit the growth of two different human breast cancer cell lines (*i.e.*, MDA-MB-435 and MCF-7). In each case, both grapefruit essence oil (OS3) and grapefruit red peel oil (OS4) extracts of the invention
 10 demonstrated the ability to inhibit the growth of each tumor cell line with greater growth inhibition being seen against the estrogen receptor positive cell line MCF-7.

In particular, grapefruit aroma (OS1), cranberry essence (OS2), grapefruit essence oil (OS3), and grapefruit red peel oil (OS4) extracts were prepared using methods described herein and by partially evaporating under nitrogen gas to remove any
 15 traces of methanol which may be to be toxic to cells. The physical characteristics of each particular extract after partial evaporation are presented in Table 3. The extracts OS1 and OS2 were soluble in DMSO, but not in water, 95% ethanol, 70% ethanol or methanol. The extracts OS3 and OS4 were soluble in 95% ethanol, but not 70% ethanol, methanol, DMSO, or water. For conducting the following experiments, stock
 20 solutions of each extract were made in DMSO (OS1 and OS2) or in 95 % ethanol (OS3 and OS4).

Table 3. Physical characteristics of four different grapefruit and cranberry extracts.

Extract	Origin	% weight loss after evaporation	Density after evaporation (g/mL)
OS1	Grapefruit aroma	32.43	0.9426
OS2	Cranberry essence	34.37	0.9820
OS3	Grapefruit essence oil	24.33	0.8382
OS4	Grapefruit red peel oil	29.21	0.8450

25 The *in vitro* assay was performed as follows. First, the human breast cancer cell lines MDA-MB435 (estrogen receptor-negative) and MCF-7 (estrogen receptor-positive) were cultured under standard conditions using, minimum essential

medium (alpha modification, 3.7 gm of sodium bicarbonate per liter, 10% v/v fetal calf serum). Media for culturing MCF-7 cells was further supplemented with 1 mM sodium pyruvate, 10 ug/mL insulin, 1% v/v fungizone (antibiotic/antimycotic, 10,000 units/mL penicillin G sodium, 10,000 ug/mL streptomycin sulfate and 25 ug/mL amphotericin B in 0.85% saline)).

Next, cells were plated at a density of 2×10^4 cells/well in 96-well, flat-bottomed tissue culture plates in a total volume of 200 uL of medium and incubated at 37°C, with or without grapefruit aroma (OS1), cranberry essence (OS2), grapefruit essence oil (OS3), or grapefruit red peel oil (OS4) extracts. The plates were incubated for 48 hours at 37°C and [3H] thymidine was then added to determine the number of dividing cells at each concentration of sample. The cells were reincubated for 4 hours, after which the medium and excess radiolabel were removed and cells were harvested and assayed for incorporated radioactivity as a measure of cell proliferation. Accordingly, the percentage of dividing cells was determined by comparing the number of disintegrations per minute of the treated cells (average of 3 wells/concentration) with that obtained for the control cells. The concentrations at which 50 % and 90 % growth inhibition occurred was determined as the IC50 and IC90 values for each extract. Results are presented in Tables 4-5 and represent the average of 3 experiments \pm SEM.

Table 4. The effect of grapefruit aroma, cranberry essence, grapefruit essence oil, and grapefruit red peel oil on the proliferation of MDA-MB-435 estrogen receptor-negative human breast cancer cells in culture.

Extract	Origin	IC50 ($\mu\text{g/mL}$)	IC90 ($\mu\text{g/mL}$)
OS1	Grapefruit aroma	>4000	>4000
OS2	Cranberry essence	>4000	>4000
OS3	Grapefruit essence oil	32.9 ± 5.2	37.0 ± 3
OS4	Grapefruit red peel oil	33.1 ± 2.5	40.0 ± 3

Table 5. The effect of grapefruit aroma, cranberry essence, grapefruit essence oil, and grapefruit red peel oil on the proliferation of MCF-7 estrogen receptor-positive human breast cancer cells in culture.

Extract	Origin	IC50 ($\mu\text{g/mL}$)	IC90 ($\mu\text{g/mL}$)
OS1	Grapefruit aroma	>4000	>4000
OS2	Cranberry essence	>4000	>4000
OS3	Grapefruit essence oil	24.1 ± 1.9	32.0 ± 4
OS4	Grapefruit red peel oil	21.0 ± 2.4	31.0 ± 1

In summary, grapefruit aroma and cranberry essence displayed no cytotoxicity against either MCF-7 (estrogen receptor positive) or MDA-MB-435 (estrogen receptor negative) human breast cancer cell lines at concentrations up to 4 mg/ml (the compounds were not tested at concentrations any higher than this due to the constraints of solubility and toxicity due to the vehicle (DMSO)). By contrast, both grapefruit essence oil and red peel oil showed high levels of anti-cancer activity against both cell lines. In particular, the dose response was extremely steep between the IC50 and IC90 values. In addition, at concentrations within 5 to 10 $\mu\text{g/ml}$ above the IC90 values, almost all of the cancerous cells were completely eliminated after incubation with grapefruit essence extract (OS3) or grapefruit red peel oil extract (OS4). These data indicate that these breast cancer cell lines are very susceptible to the type of damage inflicted by these particular extracts (or components therein), and that once a threshold amount of damage is induced, the cells undergo apoptosis.

EXAMPLE 2***IN VIVO* DEMONSTRATION OF THE ANTI-CANCER PROPERTIES OF
CRANBERRY AND GRAPEFRUIT DERIVATIVES**

The following studies were performed to examine the anti-cancer properties of
5 cranberry and grapefruit derivatives when administered to a mammal.

Four different cranberry and grapefruit extracts were prepared using methods
described herein and tested for their anti-cancer activity when administered to a test
animal having a cancer. In particular, processed grapefruit peel, concentrated pink
grapefruit juice, decharacterized cranberry, and concentrated cranberry juice were
10 tested. At least decharacterized cranberry and both the concentrated cranberry juice and
concentrated pink grapefruit juice demonstrated the ability to reduce the incidence of
tumor metastasis in a test animal with a cancer. This was taken as an indication that
certain cranberry and grapefruit derivatives of the invention (and thus, components
thereof) have powerful anticancer properties when administered to a mammal.

15 The *in vivo* assay for testing the anticancer properties of the cranberry and
grapefruit derivatives of the invention was performed as follows. First, 24 female
athymic nude mice (NCR-nu/nu), aged 3 weeks, were assigned to 1 of 5 experimental
groups and housed under standard conditions. Each group of test animals was then put
on either a control diet or test diet comprising a cranberry or grapefruit derivative. Each
20 of the diets contained 5% corn oil (wt/wt) and the amount of dextrose was adjusted to
allow for the sugar content of a particular juice being administered. The five groups of
test animals were given the following treatments: group 1, control; group 2,
concentrated cranberry juice; group 3, concentrated pink grapefruit juice; group 4,
decharacterized cranberry (5%); and group 5, processed grapefruit peel (5%);
25 After spending one week on the foregoing diets, the test animals were inoculated with
cells from the estrogen receptor-negative MDA-MB-435 human breast cancer cell line
(1×10^6 cells suspended in 50 μ L of phosphate-buffered saline (PBS)).

To carry out this procedure of experimentally seeding the test animal with a
tumorigenic cell line, mice were anesthetized with metofane, and the tumor cells were
30 injected into a right-sided mammary fat pad that had been exposed by a small incision.
The mice were weighed and the inoculation site and auxiliary lymph nodes palpated at
weekly intervals. When primary tumors became palpable, the maximum length and

width of each were measured with calipers weekly until completion of the study and corresponding surface areas were calculated. The mice were the sacrificed at 11 weeks post injection of the tumor cells.

At necropsy, body weights and primary tumor weights were recorded and the
5 number of macroscopic lung metastases was assessed using standard techniques. In
addition, the auxiliary lymph nodes and lungs were fixed in 10% neutral buffered
formalin, embedded in paraffin, sectioned, stained with hematoxylin and eosin, and
examined by light microscopy to determine the presence of metastases in these organs.
The statistically analysis of differences in the incidence of primary tumors and
10 metastases were evaluated using the chi-square test. Other statistical comparisons were
made by Student's unpaired t test with $p < 0.05$ being considered statistically significant.

In determining if any of the cranberry of grapefruit based diets influenced cancer
growth, the overall body weight and growth of mammary fat pad tumors was assessed.
The growth rate of the animals was determined to be similar in all of the test groups
15 (Fig. 1). Next, a comparison of the cumulative incidence of mammary fat pad tumors in
the different test groups, over the 11-week experimental period, was performed as
shown in Fig. 2. In groups given concentrated cranberry juice, concentrated pink
grapefruit juice, or decharacterized cranberry, the incidence of mammary fat pad tumors
was significantly reduced ($p < 0.01$). Processed grapefruit peel appeared to have little or
20 no effect on tumor incidence under these conditions. Decharacterized cranberry delayed
the onset of tumors by 4 weeks whereas concentrated cranberry juice delayed the onset
of tumors by 2 weeks.

The final incidence of mammary fat pad tumors in each dietary group was 84.7%
for the control, 78.6% for the test animals administered processed grapefruit peel, 39.4%
25 for the test animals administered concentrated cranberry juice, 38.9% for the test
animals administered concentrated pink grapefruit juice, and 32.9% for the test animals
administered decharacterized cranberry. In addition, mammary fat pad tumor size was
inhibited most significantly in test animals administered decharacterized cranberry (Fig.
4). The differences between the control and the experimental groups were statistically
30 significant except for the group given processed grapefruit peel ($p < 0.01$).

To further characterize the anticancer properties of the cranberry and grapefruit derivatives of the invention, the occurrence of lymph node and lung metastases were evaluated in test animals inoculated with a tumorigenic cell line and administered one of the above cranberry or grapefruit based test diets. The effects of administering concentrated cranberry juice, concentrated pink grapefruit juice, decharacterized cranberry, or processed grapefruit peel on lymph node metastases in a test animal are shown in Figs. 5-6. Importantly, metastases did not appear in any of the test animals administered a cranberry or grapefruit derivative until week 8. In addition, decharacterized cranberry markedly reduced the incidence of lymph node metastases when compared to the control, followed by concentrated cranberry juice and concentrated pink grapefruit juice ($p < 0.01$). No difference was observed between the group given processed grapefruit peel and control.

As yet another determination of the anticancer properties of the cranberry and grapefruit derivatives of the invention, the occurrence of macroscopically detectable lung metastases in the above test animals was also assessed (see Table 6 and Fig. 3). The incidence of lung metastases was significantly reduced in test animals given decharacterized cranberry followed by either concentrated cranberry juice or concentrated grapefruit juice, as compared to control animals ($p < 0.01$). There was no statistically significant difference between the control and the group given processed grapefruit peel.

Table 6. Effect of cranberry juice, pink grapefruit juice, decharacterized cranberry, and processed grapefruit peel on the incidence of microscopic lung metastases.

GROUP	INCIDENCE (%)	MEAN NO. PER MOUSE
control	83	4.3 ± 0.9
processed grapefruit peel	75	4.2 ± 0.7
conc. pink grapefruit juice	43	1.7 ± 0.3
conc. cranberry juice	37	1.2 ± 0.2
decharacterized cranberry	29	0.6 ± 0.1

5

IN VITRO DEMONSTRATION OF THE CHOLESTEROL LOWERING PROPERTIES OF CRANBERRY AND GRAPEFRUIT DERIVATIVES

10

15

20

30

enzyme-linked immunosorbent assay (ELISA). In particular, cells were washed and dissolved in 0.1 N NaOH for protein determination and the apo B content of the medium was calculated in μg per mg of cell protein and expressed as percent of control (medium of cells incubated with carrier liquid) and these results are presented in Table 7 and in
5 Fig. 1.

The results show that increasing concentrations of either grapefruit essence oil (OS3) or grapefruit red peel oil (OS4) extracts caused a dose-dependent reduction of apo B in the cell medium. The highest non-toxic dose of either grapefruit essence oil extract (OS3) or grapefruit red peel oil extracts (OS4) (200 $\mu\text{g}/\text{mL}$) significantly lowered
10 medium apo B by 97% and 94%, respectively. A significant apo B reduction was also produced by grapefruit essence oil extracts (OS3) at concentrations 100 and 50 $\mu\text{g}/\text{mL}$, respectively, (85% and 47%) and by grapefruit red peel oil extract (OS4) at the concentration 100 $\mu\text{g}/\text{mL}$ (66%). The concentrations required for 50% reduction of medium apo B (or IC_{50} values) were 56 $\mu\text{g}/\text{mL}$ for grapefruit essence oil extract (OS3)
15 and 70, $\mu\text{g}/\text{mL}$ for grapefruit red peel oil extract(OS4). In contrast, neither the grapefruit aroma extract (OS1) nor the cranberry essence extract (OS2) did not significantly affect levels of apo B in the medium at any of the concentrations tested.

003344:044304

Table 7. Changes in overall apo B production in HepG2 cells exposed to increasing concentrations of grapefruit extracts.

Extract	N	Conc. µg/mL	% apo B in medium
grapefruit aroma (OS1)	4	0	100 ± 20
	4	25	102 ± 15
	4	50	93 ± 23
	4	100	90 ± 22
	4	200	88 ± 30
grapefruit essence oil (OS2)	4	0	100 ± 22
	4	25	85 ± 15
	4	50	93 ± 30
	4	100	100 ± 4
	4	200	96 ± 11
grapefruit essence oil (OS3)	4	0	100 ± 6
	4	25	64 ± 30
	4	50	53 ± 5*
	4	100	15 ± 7*
	4	200	3 ± 1*
grapefruit red peel oil (OS4)	4	0	100 ± 2
	4	25	86 ± 26
	4	50	80 ± 23
	4	100	34 ± 14*
	4	200	6 ± 3*

5 **Means ± SEM. * -significantly different from control, p<0.01.**

Thus, these results show that at least two out of the four cranberry and grapefruit derivatives tested, *i.e.*, grapefruit essence oil (OS3) and grapefruit red peel oil (OS4), have cholesterol-lowering potential when tested using human liver cells (HepG2). The grapefruit essence appeared to have more potent apo B-lowering potential than the

10 grapefruit red peel oil. Moreover, using a MTT assay to assess cell viability, it was determined that none of the cranberry or grapefruit derivatives at the dosages tested

were toxic to cells. Accordingly, these extracts are healthy sources of beneficial compounds for regulated desired changes in cholesterol metabolism

EXAMPLE 4

5 ***IN VIVO* DEMONSTRATION OF THE CHOLESTEROL LOWERING PROPERTIES OF CRANBERRY AND GRAPEFRUIT DERIVATIVES**

The following studies were performed to examine the cholesterol lowering properties of cranberry and grapefruit derivatives when administered to a mammal.

In particular, four different cranberry and grapefruit extracts (*i.e.*,
10 decharacterized cranberry, concentrated (2x) cranberry juice, processed grapefruit peel, and concentrated (2x) grapefruit juice) were prepared using methods described herein and tested for their cholesterol-lowering potential using a rabbit model of hypercholesterolemia. Of the cranberry and grapefruit extracts tested, at least one extract, processed grapefruit peel, produced desired effects on cholesterol metabolism
15 when administered to a mammal.

The animal study was conducted as follows. First, rabbits were chosen as an model animal system because experimental hypercholesterolemia associated with an elevation of LDL cholesterol levels, similar to that observed in humans, can be induced by feeding the animals a low fat, cholesterol-free semipurified diet containing casein for
20 at least 3 weeks. To establish whether this effect can be counteracted by any of the cranberry or grapefruit derivatives, the animals were given casein-based diets in which these products were incorporated. After 3 weeks, cholesterolemic responses were then measured in the test animals and compared to the control animals.

The particular animals used were New Zealand White male rabbits weighing 1.6-
25 1.7 kg at the inception of the study and housed individually at constant temperature (21-24°C) and under standard light cycle conditions (12h light: 12h dark). The animals were fed ground high fiber rabbit pellets for five days after arrival and then gradually transferred (over the course of one week) to the test diets. The animals were maintained on the diets for three weeks and food and fluid consumption as well as body weight
30 changes were closely monitored.

Specifically, rabbits were assigned to five groups comprising eight animals each. The control group (CON), the group fed decharacterized cranberry (PRESS), and the group fed grapefruit pellets (PELL) was given semipurified, casein-based diets with or without respective supplements (30% each) and water to drink. The remaining two groups were given casein-based diets and either single strength cranberry juice (CRJUC) or double strength pink grapefruit juice (PGJUC) to drink. To ensure that the intake of sugars, fiber, and other components present in the administered cranberry and grapefruit derivatives did not reduce the intake of casein and other essential nutrients in the test animals, semipurified diets were modified to allow for the amounts of nutrients present in the cranberry and grapefruit derivatives. The composition of the semipurified diets alone (CON) or with cranberry or grapefruit derivative supplements is shown in Table 8. The results of nutrient analyses of the cranberry and grapefruit derivatives tested are presented in Table 9.

Table 8. Percent composition of semipurified diets used in the study.

Ingredient	Control	PRESS	PELL	CRJUC	PGJUC
Casein	26.30	37.60	37.60	31.60	30.70
Dextrose	48.20	32.70	45.10	37.90	39.60
Alfacell	13.00	11.95	--	15.60	15.20
Salt mixture (P & H)	4.00	5.70	5.70	4.80	4.67
Molasses 1:1	3.00	4.30	4.30	3.60	3.48
Vit. Mix (water sol.)	1.5	2.15	2.15	1.80	1.75
Vit. Mix (fat sol.)	0.45	0.58	0.58	0.48	0.52
Soy oil	2.22	3.10	2.75	2.67	2.58
Palm oil	1.34	1.90	1.70	1.62	1.57
Kcal/ g	3.26	3.22	3.66	3.12	3.14

PRESS - plus 30% ground decharacterized cranberry. With this supplement, the diet contained 2.28 cal/g

PELL - plus 30% ground grapefruit pellets. With this supplement, the diet contained 2.61 cal/g

CRJUC - plus cranberry juice (1x strength) instead of drinking water

PGJUC - plus pink grapefruit juice (2x normal strength) instead of drinking water

Table 9. Percent composition of cranberry and pink grapefruit products incorporated into experimental diets.

Ingredient	CR. PRESSCAKE	PGR PELLETS	CR JUICE (1x conc)	PCR JUICE (2x conc)
Protein	1.3	5.2	-	-
Sugars	0.4	-	2.73	16.4
Fiber	15.4	71.3	-	-
Minerals	0.1	10.2	-	-
Fat	0.2	1.5	-	-
Moisture	83.0	11.8	-	-
Kcal/ g	0.085	0.34	0.11	0.65

5

The growth performance for all the animal groups tested are shown in Table 10. Food consumption was comparable to the control (CON) in test animals given decharacterized cranberry (PRESS), processed grapefruit peel (PELL), and cranberry juice (CRJUC) diet but lower in animals administered the pink grapefruit juice (PGJUC) diet. Intake of casein and fiber was generally similar among all the test animals due to modifications of diet composition. Total caloric intake was significantly increased in test animals fed decharacterized cranberry (PRESS) and processed grapefruit peel (PELL) and reduced in the animals administered pink grapefruit juice (PGJUC). This, however, did not affect overall body weight gains, which remained similar, except among those animals administered the diet including cranberry juice (CRJUC).

10

15

FOOTNOTES



5 **Means \pm SEM. Statistical analysis by ANOVA plus post test by Dunnett's method.**
*** = significantly different from control, $p < 0.05$.**

These experimental results indicate that the incorporation of cranberry (PRESS),
20 cranberry juice (CRJUC), and pink grapefruit juice (PGJUC) into casein-based diet did
not counteract the elevation of serum and undesirable LDL cholesterol produced by this
diet. In contrast, incorporation of the processed grapefruit peel (PELL) significantly
reduced serum total, reduced undesirable LDL cholesterol, as well as improving the
LDL/HDL cholesterol ratio (*i.e.*, a significant decrease). In addition, a significant
25 reduction of undesirable LDL cholesterol levels, a significant increase of desirable HDL
cholesterol levels, and an improvement (*i.e.*, a significant decrease) in the LDL/HDL
cholesterol ratio, were observed in the test animals administered decharacterized

cranberry (PRESS). The test animals administer pink grapefruit juice (PGJUC) tended to have lower undesirable LDL cholesterol levels (*i.e.*, a 26% reduction) than control animals (CON) as well as improved LDL/HDL cholesterol ratios as compared to controls.. Cholesterolemic responses appeared to be largely unaffected in test animals
 5 administered cranberry juice (CRJUC).

Table 11. Serum and lipoprotein cholesterol levels in rabbits fed experimental diets.

Diet	N	Serum total cholesterol mg/dL	VLDL cholesterol mg/dL	LDL cholesterol mg/dL	HDL cholesterol mg/dL	LDL/HDL cholesterol ratio
CON	6	146 ± 12	41 ± 5	80 ± 11	30 ± 3	2.8 ± 0.5
PRESS	7	116 ± 4	22 ± 2	48 ± 3*	43 ± 1*	1.1 ± 0.1*
PELL	8	102 ± 11*	36 ± 8	30 ± 3*	33 ± 3	1.0 ± 0.2*
CRJUC	7	169 ± 12	48 ± 5	87 ± 12	37 ± 4	2.5 ± 0.5
PGJUC	7	142 ± 10	42 ± 2	59 ± 4	37 ± 4	1.7 ± 0.2*

10 Means ± SEM. Statistical analysis by ANOVA plus post test by Dunnett's method.

* = significantly different from control, p<0.05.

In summary, the above *in vivo* results demonstrated that administering processed grapefruit peel to an animal on a casein diet can counteract hypercholesterolemia
 15 induced by casein. This effect is unlikely due to merely an accompanying increase in fiber intake in these test animals because previous studies in rabbits have shown that dietary fiber does not affect casein-induced hypercholesterolemia.

Similarly, administering pink grapefruit juice into a casein diet also tended to counteract hypercholesterolemic responses induced by casein. Accordingly, it was
 20 concluded that certain grapefruit derivatives (and thus components thereof), when administered to a mammal, can improve the *in vivo* cholesterol profile of such a mammal.

2025 RELEASE UNDER E.O. 14176

EXAMPLE 5**ANALYSIS OF THE COMPONENTS OF DECHARACTERIZED CRANBERRY**

In this example, Tomah presscake (decharacterized cranberry) was determined to be enriched for a number of desirable therapeutic components. Accordingly, a detailed description of the major and minor components present in decharacterized cranberry, and suitable uses thereof, is described in the following subsections.

General Characterization of Major Components

In order to determine what major components of the decharacterized cranberry of the invention contributed to the beneficial anticancer effects when administered to a mammal (see, Example 2), a detailed analysis of the major components present in the decharacterized cranberry of the invention was performed. The decharacterized cranberry of the invention is preferably prepared using the methods described in, *e.g.*, U.S.P.N.s 5,320,861; 5,419,251. The amount of major components such as anthocyanins, phenolics, and proanthocyanidins of a decharacterized cranberry prepared using these methods (the so-called "Tomah presscake") is surprisingly enriched over the levels of these compounds found in decharacterized cranberries prepared by conventional methods (see Table 12).

Table 12. Comparison of Levels of Major Components in the Tomah Presscake as Compared to Presscake Prepared by Conventional Method

Sample	Percent Available in Fruit		
	Total Anthocyanins %	Total Phenolics %	Proanthocyanidins %
MARKHAM PRESSCAKE	6.88%	5.37%	22.72%
TOMAH PRESSCAKE	34.34%	10.52%	64.56%

Table 13. Phenolic compounds identified in decharacterized cranberry.

	Compound
1	Para-coumaric acid
2	Caffeic acid
3	Chlorogenic acid
4	Ferulic acid
5	Protocatechuic acid
6	Cinnamic acid
7	Benzoic acid
8	Gallic acid
9	Para-hydroxybenzoic acid

The biologic properties of these compounds identified as being present in the
5 cranberry are listed in a Table 15.

Analysis of the Flavanoid Compounds

The flavonoids in the decharacterized cranberry of the invention were determined using HPLC, MS, and UV spectral analysis. Standard HPLC parameters were employed and the chromatogram was monitored at 280 or 320 nm.

Using the above methods of analysis, nine different phenolic components were identified and these are listed in Table 14 below.

Table 14. Flavonoids identified in decharacterized cranberry.

	Class	Compound
1	Proanthocyanidins	polymers of flavan-3-ols
2		epicatechin oligomers
3		procyanidin B1
4		procyanidin b2
5		procyanidin b3
6	Flavan-3-ols	catechin
7		catechin gallate
8		epicatechin
9		epicatechin gallate
10		epigallocatechin gallate
11		gallocatechin gallate
12	Anthocyanins	cyanidin-3-arabinoside
13		cyanidin-3 -galactoside
14		cyanidin-3-glucoside
15		peonidin-3-arabinoside
16		peonidin-3 -galactoside
17		peonidin-3 -glucoside
18		malvidin-3 -arabinoside
19		malvidin-3 -glucoside
20	Flavonols	quercetin
21		q-3-arabinoside (avicularin)
22		q-3-galactoside (hyperin)
23		q-3-glucoside (isoquercitrin)
24		q-3-rhamnoside (quercitrin)
25		myricetin
26		m-3-arabinoside
27		m-3-rhamnoside (myricitrin)
28		m-3-digalactoside
29		kaempferol
30		isorhamnetin

T O C E H O T E T S E B O

Quercetin in particular has been shown to inhibit tumor promotion by chemical carcinogens, inactivate some enzymes involved in the metabolism of carcinogens, and inhibit LDL oxidation.

The biologic properties of this and other related compounds identified as being present in the cranberry are listed in a Table 15.

Analysis of Fiber

The fiber, i.e., pectins in the decharacterized cranberry of the invention were also determined using standard techniques. These compounds have the beneficial properties of reducing colon cancer and plasma cholesterol concentration.

Analysis of Omega-3-fatty acids and Tocotrienols

Decharacterized cranberry of the invention was also analyzed for the presence of omega-3-fatty acids and tocochromanols, a class of compounds that includes both tocopherols and tocotrienols. The method used for quantitating these compounds is based on the ability of these compounds to reduce the ferric ions (Fe^{3+}) to (Fe^{2+}). Absorption intensity is proportional to concentration, thus allowing for a determination of the amount of compound present in the sample.

Separation of individual tocotrienols was carried out using high performance liquid chromatography according to the method of Carpenter (Carpenter, Jr., A.P. *J. Amer. Oil Chemists's Soc.*, 56:668 (1979)). Detecting particular tocotrienols was conducted using UV absorption at 295 nm and identification was achieved by performing a comparison of retention times for the unknown components against known standards.

Tocotrienols in particular can inhibit growth and proliferation *in vitro* of both estrogen receptor positive or negative human breast cancer cells. The biologic properties of these compounds identified as being present in decharacterized cranberry are listed in a Table 15.

Analysis of Triterpenoids

The decharacterized cranberry of the invention was also determined to contain triterpenoids, in particular, ursolic acid. The presence of these compounds was

determined using art recognized techniques. Ursolic acid has been shown to have anti-inflammatory activity.

The biologic properties of these compounds identified as being present in the cranberry are listed in a Table 15.

5

Analysis of Other Components

The decharacterized cranberry of the invention was also determined to have several other compounds outside the compound classes mentioned above. For examples ellagic acid was also detected using art recognized techniques. This compound can stimulate apoptosis in cells and thus control the proliferation of cancer cells.

10

The biologic properties of these compounds identified as being present in the cranberry are listed in a Table 15.

Summary

15

The determinations described above reveal that the decharacterized cranberry of the invention has a remarkably high amount of anthocyanins, phenolics, and proanthocyanidins. In addition, decharacterized cranberry also contains a number of components having activity in a variety of pathways of cancer initiation, propagation, and proliferation. Accordingly, decharacterized cranberry can be considered a valuable source of therapeutic components and thus, a convenient vehicle for administering therapeutically effective mixtures of these components to a mammal, such as a human, having, *e.g.*, cancer or hypercholesterolemia. A comprehensive list of the components present in decharacterized cranberry including therapeutic uses and applications for such components, is presented in Table 15.

20

25

0034121500

Table 15.

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
CAFFEIC ACID	ABORTIFACIENT EFFECT	ANTIFERTILITY
CYANIDIN	ADENOSINE DEAMINASE INHIBITION	IMMUNODEFICIENCY
DELPHINIDIN	ANTIAPHYLACTIC ACTIVITY	ALLERGY
HYPEROSIDE	ADENOSINE A-1 RECEPTOR BINDING INHIBITION	MUSCLE ACTIVITY
KAEMPFEROL	12-HETE SYNTHESIS INHIBITION	ANALGESIC ACTIVITY
KAEMPFEROL GLYCOSIDE A	ANTIIMPLANTATION EFFECT; ANTISPERMATOGENIC EFFECT	IMMUNOLOGIC; FERTILITY
KAEMPFEROL GLYCOSIDE B	ANTIIMPLANTATION EFFECT	IMMUNOLOGIC
KAEMPFEROL TETRAACETATE	PLATELET AGGREGATION INHIBITION	PLATELET AGGREGATION
KAEMPFEROL TRIMETHYL ETHER	ANTIMUTAGENIC ACTIVITY	ANTIMUTAGENIC ACTIVITY
KAEMPFEROL, 3-4'-7-TRIMETHYL	CYCLIC AMP INHIBITION	ANTIINFLAMMATORY
KAEMPFEROL, 3-4'-7-TRI-O-METHYL	COMPLEMENT CLASSICAL PATHWAY INHIBITION; COMPLEMENT-ALTERNATE PATHWAY STIMULATION	IMMUNITY
KAEMPFEROL, 3-4'-DI-O-METHYL	COMPLEMENT CLASSICAL PATHWAY INHIBITION; COMPLEMENT-ALTERNATE PATHWAY STIMULATION	IMMUNITY
KAEMPFEROL, 3-O-METHYL	CYTOTOXIC ACTIVITY	ANTICANCER
KAEMPFEROL, 3-RUTINOSYL	RADICAL SCAVENGING EFFECT	ANTIOXIDANT
KAEMPFEROL, 4'-5-7-TRI-O-METHYL: (FLAVONOID)	ANTIBACTERIAL ACTIVITY; ANTIMYCOBACTERIAL ACTIVITY; PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIBACTERIAL ACTIVITY; ANTIMYCOBACTERIAL ACTIVITY; ANTIINFLAMMATORY
KAEMPFEROL, 6-8-DI-C-METHYL: 3-METHYL ETHER (FLAVONOID)	PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIINFLAMMATORY
KAEMPFEROL, 6-HYDROXY: (FLAVONOID)	DNA POLYMERASE I INHIBITION; INOTROPIC EFFECT POSITIVE; REVERSE TRANSCRIPTASE INHIBITION	ANTITUMOR; MUSCLE; ANTICANCER
KAEMPFEROL, 6-HYDROXY: 3-6-7-TRIGLUCOSIDE (FLAVONOID)	INOTROPIC EFFECT POSITIVE	MUSCLE
KAEMPFEROL, 6-HYDROXY: 3-6-DI-GLUCOSIDE (FLAVONOID)	INOTROPIC EFFECT POSITIVE	MUSCLE
KAEMPFEROL, 6-HYDROXY: 3-O-BETA-D-GLUCOSIDE (FLAVONOID)	ATP-ASE(Na+/K+) INHIBITION; INOTROPIC EFFECT POSITIVE	BLOOD; MUSCLE
KAEMPFEROL, 6-HYDROXY: 3-RUTINOSYL-6-GLUCOSIDE (FLAVONOID)	INOTROPIC EFFECT POSITIVE	MUSCLE
KAEMPFEROL, 6-HYDROXY: 4'-METHYL ETHER 3-7-DIHRHAMNOSIDE (FLAVONOID)	CYTOTOXIC ACTIVITY	ANTICANCER
KAEMPFEROL, 6-HYDROXY-3-7-DIMETHYL ETHER (FLAVONOID)	SMOOTH MUSCLE RELAXANT ACTIVITY; SPASMOLYTIC ACTIVITY	MUSCLE; NERVOUS SYSTEM
KAEMPFEROL, 6-METHOXY: (FLAVONOID)	ANTICYTOTOXIC ACTIVITY; ATP-ASE (Ca++) INHIBITION; BETA-GLUCURONIDASE INHIBITION; CYTOTOXIC ACTIVITY; HISTAMINE	ANTICANCER; BLOOD; DETOXIFICATION; ANTICANCER; ANTIOXIDANT

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

	RELEASE INHIBITION	
KAEMPFEROL, 6-METHOXY: 3-7-DIMETHYL ETHER (FLAVONOID)	ANTIVIRAL ACTIVITY	ANTIVIRAL ACTIVITY
KAEMPFEROL, 6-METHOXY: 3-METHYL ETHER (FLAVONOID)	ANTIVIRAL ACTIVITY	ANTIVIRAL ACTIVITY
KAEMPFEROL, 6-METHOXY: 3-O-BETA-G-(1-6)-ROBINOSIDE (FLAVONOID)	CALCIUM ION RELEASE INHIBITION; CYTOTOXIC ACTIVITY	BLOOD; ANTICANCER
KAEMPFEROL, 6-METHOXY: 3-O-BETA-D-ROBINOSIDE (FLAVONOID)	ALDOL REDUCTASE (LENS) INHIBITION	ANTINEOPLASTIC
KAEMPFEROL-3-(2-3-DIACETOXY-4-PARA-COUMAROYL)-RHOMNOSIDE (FLAVONOID)	ANTIFUNGAL ACTIVITY	ANTIFUNGAL ACTIVITY
KAEMPFEROL-3-4'-7-TRIMETHYL ETHER (FLAVONOID)	ANTICRUSTACEAN ACTIVITY; ANTIVIRAL ACTIVITY; CYTOTOXIC ACTIVITY	ANTICRUSTACEAN ACTIVITY; ANTIVIRAL ACTIVITY; ANTICANCER
KAEMPFEROL-3-4'-DIMETHYL ETHER (FLAVONOID)	ANTIVIRAL ACTIVITY; REVERSE TRANSCRIPTASE INHIBITION	ANTIVIRAL ACTIVITY; ANTICANCER
KAEMPFEROL-3-7-DIMETHYL ETHER (FLAVONOID)	ANTIVIRAL ACTIVITY	ANTIVIRAL ACTIVITY
KAEMPFEROL-3-7-DI-O-BETA-D-GLUCOSIDE (FLAVONOID)	CITRATE SYNTHASE INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-BETA-DIRHAMNOSIDE (FLAVONOID)	PLANT ROOT GROWTH INHIBITION	PLANT GROWTH

TDETHD: TETSECO

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
KAEMPFEROL-3-GLUCOSIDE-7-RHAMNOSIDE (FLAVONOID)	ANTIOXIDANT ACTIVITY; CYCLOOXYGENASE STIMULATION; LIPOXYGENASE INHIBITION; PLATELET AGGREGATION INHIBITION	ANTIOXIDANT; ANTIINFLAMMATORY; ANTIOXIDANT; PLATELET AGGREGATION
KAEMPFEROL-3-NEOHESPERIDOSIDE (FLAVONOID)	BRONCHODILATOR ACTIVITY	LUNG
KAEMPFEROL-3-O-(2-4-(DI-PARA-COUMAROYL)-BETA-D-GLUCOSIDE) (FLAVONOID)	ANTIINFLAMMATORY ACTIVITY	ANTIINFLAMMATORY ACTIVITY
KAEMPFEROL-3-O-(2"-6"-DI-O-TRANS-PARA-COUMAROYL-BETA-D-GLUCOSIDE) (FLAVONOID)	REPELLENT ACTIVITY (MOLLUSK)	MOLLUSK
KAEMPFEROL-3-O-(2"-O-GALLOYL)-GLUCOSIDE (FLAVONOID)	ANGIOTENSIN-CONVERTING ENZYME INHIBITION	VASOPRESSOR
KAEMPFEROL-3-O-(6"-ACETYL)-BETA-D-GLUCOSIDE (FLAVONOID)	CHROMOSOME ABERRATION INHIBITION	ANTICANCER
KAEMPFEROL-3-O-(6"-O-ACETYL)-BETA-D-GLUCOSIDE (FLAVONOID)	GLUCOSYLTRANSFERASE INHIBITION	LIVER
KAEMPFEROL-3-O-(6"-O-ACETYL)-GLUCOSIDE (FLAVONOID)	CYTOTOXIC ACTIVITY	ANTICANCER
KAEMPFEROL-3-O-(6"-O-TRANS-PARA-COUMAROYL)-BETA-D-GLUCOPYRANOSIDE (FLAVONOID)-	ANTIVIRAL ACTIVITY; CYTOTOXIC ACTIVITY	ANTIVIRAL ACTIVITY; ANTICANCER
KAEMPFEROL-3-O-[2-O-ALPHA-L-RHAMNOPYRANOSYL-4-O-BETA-D-GLUCOPYRANOSYL]-BETA	PLATELET AGGREGATION INHIBITION	PLATELET AGGREGATION
KAEMPFEROL-3-O-ALPHA-(6"-PARAPCOUMAROYL-GLUCOSYL-BETA-1-2-RHAMNOSIDE) (FLAVONOID)	FEEDING DETERRENT (INSECT)	INSECTICIDAL
KAEMPFEROL-3-O-ALPHA-ARABINOPYRANOSIDE (FLAVONOID)	PROTEIN TYROSINE KINASE INHIBITION	VASCULAR
KAEMPFEROL-3-O-ALPHA-L-(2"-3"-DI-TRANS-PARA-COUMAROYL)-RHAMNOPYRANOSIDE (FLAVONOID)	ANTIBACTERIAL ACTIVITY; ANTIBACTERIAL ACTIVITY; CYTOTOXIC ACTIVITY	ANTIBACTERIAL ACTIVITY; ANTIBACTERIAL ACTIVITY; ANTICANCER
KAEMPFEROL-3-O-ALPHA-L-ARABINOFURANOSYL-7-O-ALPHA-L-RHAMNOPYRANOSIDE (FLAVONOID)	ANTIBACTERIAL ACTIVITY; ANTIYEAST ACTIVITY	ANTIBACTERIAL ACTIVITY; ANTIYEAST
KAEMPFEROL-3-O-ALPHA-L-RHAMNOPYRANOSYL-(1-6)-BETA-D-GLUCOPYRANOSIDE (FLAVONOID)	ANALGESIC ACTIVITY	ANALGESIC ACTIVITY
KAEMPFEROL-3-O-ARABINOPYRANOSIDE-2"-GALLATE (FLAVONOID)	ANTIOXIDANT ACTIVITY	ANTIOXIDANT

KAEMPFEROL-3-O-BETA-D-GLUCOPYRANOSYL(1-2)-(ALPHA-L-RHAMNOPYRANSOYL(1-6))-BETA	ANTIBACTERIAL ACTIVITY	ANTIBACTERIAL ACTIVITY
KAEMPFEROL-3-O-BETA-D-GLUCOSIDE-7-RHAMNOSIDE (FLAVONOID)	VASODILATOR ACTIVITY	VASCULAR
KAEMPFEROL-3-O-BETA-D-NEOHESPERIDOSIDE (FLAVONOID)	PLATELET AGGREGATION INHIBITION	PLATELET AGGREGATION
KAEMPFEROL-3-O-BETA-D-RUTINOSIDE (FLAVONOID)	XANTHINE OXIDASE INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-O-BETA-D-SOPHOROSIDE (FLAVONOID)	ANALGESIC ACTIVITY; ANTIINFLAMMATORY ACTIVITY; BRONCHODILATOR ACTIVITY	ANALGESIC ACTIVITY; ANTIINFLAMMATORY ACTIVITY; LUNG
KAEMPFEROL-3-O-BETA-D-XYLOFURANOSIDE (FLAVONOID)	LIPID PEROXIDE FORMATION INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-O-BETA-D-XYLOFURANOSYL(1-2)-GALACTOPYRANOSIDE (FLAVONOID)	LIPID PEROXIDE FORMATION INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-O-BETA-D-XYLOFURANOSYL(1-2)-BETA-D-ARABINOPYRANOSIDE (FLAVONOID)	EXOCYTOSIS STIMULATION	ANTICANCER
KAEMPFEROL-3-O-METHYL ETHER (FLAVONOID)	XANTHINE OXIDASE INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-O-NEOHESPERIDOSIDE (FLAVONOID)	ANTIBACTERIAL ACTIVITY; ANTIOXIDANT ACTIVITY; ANTIYEAST ACTIVITY	ANTIBACTERIAL ACTIVITY; ANTIOXIDANT; ANTIYEAST

10/10/2023 10:10:10 AM

Table 15 continued

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
KAEMPFEROL-3-O-RHAMNOSIDE (FLAVONOID)	COMPLEMENT ALTERNATEIVE PATHWAY INHIBITION; COMPLEMENT CLASSICAL PATHWAY INHIBITION; COMPLEMENT ALTERNATE PATHWAY STIMULATION; LIPID PEROXIDE FORMATION INHIBITION	IMMUNITY; ANTIOXIDANT
KAEMPFEROL-3-O-RHAMNOSYL- DIGLUCOSIDE (FLAVONOID)	ANTIPRURITIC ACTIVITY	ALLERGY
KAEMPFEROL-3-O-RUTINOSIDE (FLAVONOID)	ADENOSINE A-1 RECEPTOR BINDING INHIBITION; ANTIBACTERIAL ACTIVITY; ANTIYEAST ACTIVITY; CHRONOTROPIC EFFECT NEGATIVE; COMPLEMENT ALTERNATIVE PATHWAY INHIBITION; COMPLEMENT-ALTERNATE PATHWAY STIMULATION; EPSTEIN-BARR VIRUS EARLY ANTIGEN ACTIVATION INHIBITION; HYPOTENSIVE ACTIVITY; SMOOTH MUSCLE RELAXANT ACTIVITY; SPASMOLYTIC ATIVITY	MUSCLE ACTIVITY; ANTIBACTERIAL ACTIVITY; ANTIYEAST; HEART; IMMUNITY; IMMUNITY; CANCER; HYPERTENSION; MUSCLE; NERVOUS SYSTEM
KAEMPFEROL-3-O-SOPHOROSIDE (FLAVONOID)	LIPID PEROXIDE FORMATION INHIBITION	ANTIOXIDANT
KAEMPFEROL-3-RHAMNOSIDE (FLAVONOID)	ANTIOXIDANT ACTIVITY; CYCLOOXYGENASE STIMULATION; LIPOXYGENASE INHIBITION	ANTIOXIDANT; ANTIINFLAMMATORY; ANTIOXIDANT
KAEMPFEROL-3-O-RHAMNOSYL-7- GLUOSIDE (FLAVONOID)	ANTIMUTAGENIC ACTIVITY	ANTIMUTAGENIC ACTIVITY
KAEMPFEROL-3-RUTINOSIDE (FLAVONOID)	ANTICYTOTOXIC ACTIVITY; AUTOPHAGY STIMULATION; CYTOTOXIC ACTIVITY; INOTROPIC EFFECT POSITIVE	ANTICANCER; IMMUNOLOGY; ANTICANCER; MUSCLE
KAEMPFEROL-3-SULFATE (FLAVONOID)	GLUCOSE TRANSPORT STIMULATION	DIABETES
KAEMPFEROL-4'-O-BETA-D- GLUCOSIDE (FLAVONOID)	HYPERTENSIVE ACTIVITY	HYPERTENSION
KAEMPFEROL-4'-O-METHYL ETHER (FLAVONOID)	ANTICHOLINERGIC ACTIVITY; ANTIINFLAMMATORY ACTIVITY; BARBITURATE POTENTIATION; CHRONOTROPIC EFFECT POSITIVE	NERVOUS SYSTEM; ANTIINFLAMMATORY ACTIVITY; DRUG; HEART
MALVIDIN	ABORTIFACIENT EFFECT	ANTIFERTILITY
MYRICETIN	12-HETE SYNTHESIS INHIBITION	ANALGESIC ACTIVITY
MYRICETIN	PLANT ROOT GROWTH INHIBITION	PLANT GROWTH
MYRICENTIN RHAMNOSIDE	ANTIULCER ACTIVITY; SMOOTH MUSCLE RELAXANT ACTIVITY	ANTIULCER; MUSCLE
MYRICETIN TRIMETHYL ETHER	12-HETE SYNTHESIS INHIBITION; SPASMOLYTIC ACTIVITY	ANALGESIC ACTIVITY; NERVOUS SYSTEM
MYRICETIN-3-ARABINOSIDE	ANTIOXIDNT ACTIVITY; CYCLOOXYGENASE STIMULATION; LIPOSYGENASE INHIBITION	ANTIOXIDANT; ANTIINFLAMMATORY; ANTIOXIDANT



Table 15 continued

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
PELARGONIDIN-5-7-DIMETHYL ETHER 3-O-ALPHA-L-RHAMNOSIDE	ANTIHYPERGLYCEMIC ACTIVITY	DIABETES
PROANTHOCYANIDIN	ANTIFUNGAL ACTIVITY; CYTOTOXIC ACTIVITY	ANTIFUNGAL ACTIVITY; ANTICANCER
PROCYANIDIN	ANTIFUNGAL ACTIVITY	ANTIFUNGAL ACTIVITY
QUERCETIN	12-HETE SYNTHESIS INHIBITION	ANALGESIC ACTIVITY
QUERCETIN,4'-7-DIMETHYL	XANTHINE OXIDASE INHIBITION	ANTIOXIDANT
QUERCETIN,4'-7-DI-O-BENZOYL	ANTIVIRAL ACTIVITY (PLANT PATHOGENS)	ANTIVIRAL ACTIVITY
QUERCETIN,4'-7-DI-O-METHYL	MUTAGENIC ACTIVITY	ANTICANCER
QUERCETIN,5-7-DI-O-METHYL	MUTAGENIC ACTIVITY	ANTICANCER
QUERCETIN,5-O-METHYL	MUTAGENIC ACTIVITY; VIRAL INFECTIVITY INHIBITION	ANTICANCER; INFECTION
QUERCETIN,6-8-DI-C-METHYL: 3-7- DIMETHYL ETHER	PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIINFLAMMATORY
QUERCETIN,6-8-DI-C-METHYL: 3- METHYL ETHER	ANTIBACTERIAL ACTIVITY; PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIBACTERIAL ACTIVITY; ANTIINFLAMMATORY
QUERCETIN,6-C-METHYL: 3-3'-7- TRIMETHYL ETHER	PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIINFLAMMATORY
QUERCETIN,6-C-METHYL: 3-7- DIMETHYL ETHER	ANTIBACTERIAL ACTIVITY; PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIBACTERIAL ACTIVITY; ANTIINFLAMMATORY
QUERCETIN,6-C-METHYL: 3-METHYL ETHER	ANTIBACTERIAL ACTIVITY; PROSTAGLANDIN SYNTHESIS INHIBITION	ANTIBACTERIAL ACTIVITY; ANTIINFLAMMATORY
QUERCETIN,7-METHOXY:	ANTIMUTAGENIC ACTIVITY	ANTIMUTAGENIC ACTIVITY
QUERCETIN,7-O-BENZOYL:	CYCLIC GMP INHIBITION; IMMUNOPRECIPITATION REACTION	ANTIINFLAMMATORY; IMMUNOLOGY
QUERCETIN, DIHYDRO: (DL)	ADENOSINE RECEPTOR BINDING INHIBITION	MUSCLE ACTIVITY
QUERCETIN, DIHYDRO: 3-O- ACETATE (FLAVONOID)	SWEETENING EFFECT	TASTE
QUERCETIN, ISO:	REVERSE TRANSCRIPTASE INHIBITION; SPASMOLYTIC ACTIVITY	ANTICANCER; NERVOUS SYSTEM
QUERCETIN,METHYL: (FLAVONOID)	RNA SYNTHESIS INHIBITION	ANTICANCER
QUERCETIN, PENTAACETYL: (FLAVONOID)	PHOSPHODIESTERASE INHIBITION	ANTIINFLAMMATORY
QUERCETIN, PENTAHYDROXY- ETHYL: (FLAVONOID)	CELL MEMBRANE PERMEATION	ANTICANCER
QUERCETIN, PENTAMETHOXY: (FLAVONOID)	ANTIMUTAGENIC ACTIVITY	ANTIMUTAGENIC ACTIVITY
QUERCETIN, PENTAMETHYL: (FLAVONOID)	ANTIVIRAL ACTIVITY; MUTAGENIC ACTIVITY; PHOSPHODIESTERASE INHIBITION; SPASMOLYTIC ACTIVITY; SPASMOLYTIC ACTIVITY	ANTIVIRAL ACTIVITY; ANTICANCER; ANTIINFLAMMATORY; NERVOUS SYSTEM; NERVOUS SYSTEM

P.03 E.140 : 12 E.140

Table 15 continued

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
QUERCETIN-3-4'-DI-O-BETA-D-GLUCOSIDE	CATALASE INHIBITION; CELL PROLIFERATION INHIBITION; GLUTAMATE-OXALOACETATE-TRANSAMINASE INHIBITION; GLUTAMATE-OXALOACETATE-TRANSAMINASE STIMULATION; IMMUNOSTIMULANT ACTIVITY; NADPH-CYTOCHROME C REDUCTASE STIMULATION; NADPH-CYTOCHROME C REDUCTASE STIMULATION	ANTIOXIDANT; ANTICANCER; HEPATOPROTECTION; HEPATOPROTECTION; IMMUNOLOGY; ANTICANCER; ANTICANCER
QUERCETIN-3-7-DIGLUCOSIDE (FLAVONOID)	ANTIOXIDANT ACTIVITY; ATP-ASE (NA+/K+) INHIBITION; CYCLOOXYGENASE STIMULATION; INOTROPIC EFFECT POSITIVE; LIPOXYGENASE INHIBITION	ANTIOXIDANT; BLOOD; ANTIINFLAMMATORY; MUSCLE; ANTIOXIDANT
QUERCETIN-3-ACETATE, DIHYDRO: (FLAVONOID)	MUTAGENIC ACTIVITY; SWEETENING EFFECT	ANTICANCER; TASTE
QUERCETIN-3-ALPHA-ARABINOPYRANOSIDE-2"-GALLATE (FLAVONOID)	ANTIBACTERIAL ACTIVITY	ANTIBACTERIAL ACTIVITY
QUERCETIN-3-GENTIOBIOOSIDE (FLAVONOID)	ANTIOXIDANT ACTIVITY	ANTIOXIDANT
QUERCETIN-3-GLUCOSIDE-7-RHAMNOSIDE	ANTIOXIDANT ACTIVITY; CYCLOOXYGENASE STIMULATION; LIPOXYGENASE INHIBITION	ANTIOXIDANT; ANTIINFLAMMATORY; ANTIOXIDANT
QUERCETIN-3-METHYL ETHER (FLAVONOID)	ALDOL REDUCTASE (LENS) INHIBITION; ANTIANAPHYLACTIC ACTIVITY; ANTIBACTERIAL ACTIVITY; ANTIFUNGAL ACTIVITY; ANTIOXIDANT ACTIVITY; ANTIVIRAL ACTIVITY; HISTAMINE RELEASE INHIBITION; SUPEROXIDE RADICAL SCAVENGING ACTIVITY	ANTINEOPLASTIC; ALLERGY; ANTIBACTERIAL ACTIVITY; ANTIFUNGAL ACTIVITY; ANTIOXIDANT; ANTIVIRAL ACTIVITY; ANTIOXIDANT; ANTIOXIDANT
QUERCETIN-3-O-(2"-6"-ALPHA-L-DIRHAMNOPYRONOSYL)-BETA-D-GALACTOPYRANOSIDE (FLAVONOID)	OVIPOSITION STIMULATION	FERTILITY
QUERCETIN-3-O-(2"-6"-O-DIGALLOYL)-BETA-D-GALACTOPYRANOSIDE (FLAVONOID)	HIV-1 INTEGRASE INHIBITION	AIDS
QUERCETIN-3-O-(2-BETA-D-XYLOPRANOSYL-RUTINOSIDE) (FLAVONOID)	OVIPOSITION INHIBITION	FERTILITY
QUERCETIN-3-O-(2"-GALLOYL)-ALPHA-L-ARABINOPYRANOSIDE (FLAVONOID)	HIV-1 INTEGRASE INHIBITION	AIDS
QUERCETIN-3-O-(2"-GALLOYL)-	HIV-1 INTEGRASE INHIBITION	AIDS

BETA-D-GALACTOPYRANOSIDE (FLAVONOID)		
QUERCETIN-3-O-(2"-GALLOYL)- BETA-D-GLUCOSIDE (FLAVONOID)	HEMOLYSIS INHIBITORY ACTIVITY	BLOOD
QUERCETIN-3-O-(2"-O-BALLOYL)- BETA-D-GALACTOSIDE (FLAVONOID)	CYTOTOXIC ACTIVITY	ANTICANCER
QUERCETIN-3-O-(2"-O-GALLOYL)- GLUCOSIDE (FLAVONOID)	ANGIOTENSIN-CONVERTING ENZYME INHIBITION; MOLLUSCICIDAL ACTIVITY	VASOPRESSOR; MOLLUSCICIDAL ACTIVITY
QUERCETIN-3-O-(6"-GALLOYL)- BETA-D-GLUCOSIDE (FLAVONOID)	HEMOLYSIS INHIBITORY ACTIVITY	BLOOD
QUERCETIN-3-O-(6"-GALLOYL)- GLUCOSIDE (FLAVONOID)	SEROTONIN ANTAGONIST ACTIVITY; SPASMOGENIC ACTIVITY	BRAIN; NERVOUS SYSTEM
QUERCETIN-3-O-(6"-O-ACETYL)- BETA-D-GLUCOSIDE (FLAVONOID)	GLUCOSYLTRANSFERASE INHIBITION; PLATELET AGGREGATION INHIBITION	LIVER; PLATELET AGGREGATION
QUERCETIN-3-O-ALPHA-(6"-PARA- COUMAROYL-GLUCOSYL-BETA-1-2- RHAMNOSIDE) (FLAVONOID)	FEEDING DETERRENT (INSECT)	INSECTICIDAL
QUERCETIN-3-O-ALPHA-L- ARABINOPYRANOSIDE (FLAVONOID)	REVERSE TRANSCRIPTASE INHIBITION	ANTICANCER
QUERCETIN-3-O-ALPHA-I- ARABINOPYRANOSYL(1-2)-ALPHA- L-RHAMNOPYRANOSIDE (FLAVONOID)	ANTIALLERGENIC ACTIVITY	ALLERGY
QUERCETIN-3-O- ARABINOFURANOSIDE (FLAVONOID)	CYTOTOXIC ACTIVITY; HISTAMINE RELEASE INHIBITION	ANTICANCER; ANTIOXIDANT

2025-04-10 14:34:00

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
QUERCETIN-3-O-BETA-D-FALACTOSE(1-6)-O-ALPHA-L-RHAMNOSE (FLAVONOID)	ANTIBACTERIAL ACTIVITY; TOPOISOMERASE II INHIBITION	ANTIBACTERIAL ACTIVITY; ANTICANCER
QUERCETIN-3-O-BETA-D-GALACTOSIDE-2"-GALLATE (FLAVONOID)	PLATELET AGGREGATION INHIBITION	PLATELET AGGREGATION
QUERCETIN-3-O-BETA-D-GALACTURONIDE (FLAVONOID)	INTESTINAL MOTILITY INHIBITION	DIGESTION
QUERCETIN-3-O-BETA-D-GLUCOPYRANOSIDE-2"-GALLATE (FLAVONOID)	MOLLUSCICIDAL ACTIVITY	MOLLUSCICIDAL ACTIVITY
QUERCETIN-3-O-BETA-D-GLUCOPYRANOSYL(1-2)-BETA-D-GLUCOPYRANOSIDE (FLAVONOID)	ANTIBACTERIAL ACTIVITY	ANTIBACTERIAL ACTIVITY
QUERCETIN-3-O-BETA-D-GLUCOPYRANOSYL(1-6)-BETA-D-GALACTOPYRANOSIDE (FLAVONOID)	OVIPOSITION STIMULATION	FERTILITY
QUERCETIN-3-O-BETA-D-GLUCORONIDE (FLAVONOID)	ANGIOTENSIN-CONVERTING ENZYME INHIBITION; INTESTINAL MOTILITY INHIBITION	VASOPRESSOR; DIGESTION
QUERCETIN-3'-O-BETA-D-GLUCOSIDE	ANTIBACTERIAL ACTIVITY; ANTIULCER ACTIVITY; CITRATE SYNTHASE INHIBITION; SMOOTH MUSCLE RELAXANT ACTIVITY	ANTIBACTERIAL ACTIVITY; ANTIULCER; ANTIOXIDANT; MUSCLE
QUERCETIN-3-O-BETA-D-GLUCURONOPYRANOSIDE (FLAVONOID)	ANTIHEPATOTOXIC ACTIVITY; GLUTAMATE-PYRUVATE-TRANSAMINASE INHIBITION; GLUTATHIONE-S-TRANSFERASE INDUCTION	ANTIHEPATOTOXIC ACTIVITY; HEPATOPROTECTION; ANTICANCER
QUERCETIN-3-O-BETA-D-XYLOPYRANOSIDE (FLAVONOID)	REVERSE TRANSCRIPTASE INHIBITION	ANTICANCER
QUERCETIN-3-O-BETA-D-SYLOPYRANOSYL(1-2)-BETA-D-GLUCOPYRANOSIDE (FLAVONOID)	CHROMOSOME ABERRATION INHIBITION	ANTICANCER
QUERCETIN-D-O-BETA-GLUCURONIDE (FLAVONOID)	CYTOTOXIC ACTIVITY	ANTICANCER
QUERCETIN-3-O-D-ROBINOSIDE (FLAVONOID)	ANTIOXIDANT ACTIVITY	ANTIOXIDANT
QUERCETIN-3-O-GALACTOSIDE OCTAACETATE	PLATELET AGGREGATION INHIBITION	PLATELET AGGREGATION
QUERCETIN-3-O-GALACTOSYL-(1-6)-GLUCOSIDE (FLAVONOID)	ANTIBACTERIAL ACTIVITY	ANTIBACTERIAL ACTIVITY
QUERCETIN-3-O-GENTIOBIOSIDE (FLAVONOID)	SPASMOLYTIC ACTIVITY	NERVOUS SYSTEM
QUERCETIN-3-O-GLUCO-RHAMNO-ARABINOSIDE (FLAVONOID)	ANTIINFLAMMATORY ACTIVITY	ANTIINFLAMMATORY ACTIVITY

Table 15 continued

COMPOUND	BIOLOGICAL ACTIVITY	UTILITY
QUERCETIN-3-SULFATE	ANTIOXIDANT ACTIVITY; SUPEROXIDE RADICAL SCAVENGING ACTIVITY	ANTIOXIDANT
QUERCETIN-4'-7-DIMETHYL ETHER	ANTIVIRAL ACTIVITY(PLANT PATHOGENS); COMPLEMENT ALTERNATIVE PATHWAY INHIBITION; COMPLEMENT CLASSICAL PATHWAY, INHIBITION	ANTIVIRAL ACTIVITY; IMMUNITY
QUERCETIN-4'-GLUCOSIDE	ANTIOXIDANT ACTIVITY; LIPID PEROXIDE FORMATION INHIBITION; QUINONE REDUCTASE INDUCTION	ANTIOXIDANT; ANTICANCER
QUERCETIN-4'-O-BETA-GLUCOPYRANOSIDE-6"-GALLATE	ANTIOXIDANT ACTIVITY	ANTIOXIDANT
QUERCETIN-4'-O-GLUCOPYRANOSYL-6"-GALLATE	ANTIOXIDANT ACTIVITY	ANTIOXIDANT
QUERCETIN-6-METHOXY: 3-3'-DIMETHYL ETHER	ANTIVIRAL ACTIVITY	ANTIVIRAL ACTIVITY
QUERCETIN-ACETYL-TRISULFATE	ALCOHOL DEHYDROGENASE INHIBITION; LACTATE DEHYDROGENASE INHIBITION; MALATE DEHYDROGENASE INHIBITION	HYPERTENSION; ANTIOXIDANT
QUERCETIN-RHAMNOSIDE	ANTIULCER ACTIVITY; SMOOTH MUSCLE RELAXANT ACTIVITY	ANTIULCER; MUSCLE
QUERCITRIN	ADENOSINE A-1 RECEPTOR BINDING INHIBITION; ALCOHOL DEHYDROGENASE INHIBITION; ALDOL REDUCTASE (LENS) INHIBITION	MUSCLE ACTIVITY; HYPERTENSION; ANTINEOPLASTIC
TANIN	ANTIBACTERIAL ACTIVITY; ANTIFUNGAL ACTIVITY; ANTIHCG ACTIVITY; ANTIHISTAMINE ACTIVITY; ANTIINFLAMMATORY ACTIVITY; ANTIOXIDANT ACTIVITY; ANTI-PREGNANT MARE SERUM GONADOTROPIN ACTIVITY; ANTITUMOR ACTIVITY; CALCIUM ION UPTAKE STIMULATION; CYTOTOXIC ACTIVITY; ESTROUS CYCLE DISRUPTION EFFECT; FEEDING DETERRENT (INSECT); FEEDING SSTIMULANT (INSECT); HEPATOTOXIC ACTIVITY; NO-SYNTHASE INHIBITION; ORNITHINE DECARBOXYLASE INHIBITION; POLYGLYCOHYDROLASE INHIBITION; UTERINE STIMULANT EFFECT	HORMONE; ANTIBACTERIAL ACTIVITY; ANTIFUNGAL ACTIVITY; HORMONE; ALLERGY; ANTIINFLAMMATORY ACTIVITY; ANTIOXIDANT; HORMONE; ANTITUMOR ACTIVITY; BLOOD; ANTICANCER; FERTILITY; INSECTICIDAL; LIVER; ANTITUMOR; IMMUNITY; FERTILITY
URSOLIC ACID	ACID PHOSPHATASE INHIBITION; ADENOSINE DEAMINASE INHIBITION	ANTINEOPLASTIC; IMMUNODEFICIENCY

T06410: 12458350

